

Modulating exercise-induced hormesis

Does less equal more?

Peake, Jonathan M.; Markworth, James F.; Nosaka, Kazunori; Raastad, Truls; Wadley, Glenn D.; Coffey, Vernon G.

Published in:
Journal of Applied Physiology

DOI:
[10.1152/jappphysiol.01055.2014](https://doi.org/10.1152/jappphysiol.01055.2014)

Published: 01/08/2015

Document Version:
Peer reviewed version

[Link to publication in Bond University research repository.](#)

Recommended citation(APA):

Peake, J. M., Markworth, J. F., Nosaka, K., Raastad, T., Wadley, G. D., & Coffey, V. G. (2015). Modulating exercise-induced hormesis: Does less equal more? *Journal of Applied Physiology*, 119(3), 172-189. <https://doi.org/10.1152/jappphysiol.01055.2014>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

For more information, or if you believe that this document breaches copyright, please contact the Bond University research repository coordinator.

1 JAPPL-01055-2014 R1

2 **Modulating exercise-induced hormesis: does less equal more?**

3 Running title: Exercise-induced hormesis

4 Jonathan M. Peake^{1,2}, James F. Markworth³, Kazunori Nosaka⁴, Truls Raastad⁵, Glenn
5 Wadley⁶, Vernon Coffey^{7,8}

6

7 ¹ School of Biomedical Sciences and Institute of Health and Biomedical Innovation,
8 Queensland University of Technology, Brisbane, Australia

9 ² Centre of Excellence for Applied Sports Science Research, Queensland Academy of Sport,
10 Brisbane, Australia

11 ³ Liggins Institute, University of Auckland, Auckland, New Zealand

⁴ School of Exercise and Health Sciences, Centre for Exercise and Sports Science Research,
Edith Cowan University, Joondalup, Australia

⁵ Norwegian School of Sport Sciences, Oslo, Norway

⁶ School of Exercise and Nutrition Sciences, Center for Physical Activity and Nutrition
Research, Deakin University, Melbourne, Australia

12 ⁷ School of Exercise and Nutrition Sciences and Institute of Health and Biomedical
13 Innovation, Queensland University of Technology, Brisbane, Australia

14 ⁸ Bond Institute of Health and Sport and Faculty of Health Sciences and Medicine, Bond
15 University, Gold Coast, Australia

Corresponding author:

Jonathan Peake
Institute for Health and Biomedical Innovation
Kelvin Grove, QLD 4059
Brisbane, Australia
Email: jonathan.peake@qut.edu.au
Phone: +61 7 3138 6140

ABSTRACT

16 Hormesis encompasses the notion that low levels of stress stimulate or upregulate
17 existing cellular and molecular pathways that improve the capacity of cells and organisms to
18 withstand greater stress. This notion underlies much of what we know about how exercise
19 conditions the body and induces long-term adaptations. During exercise, the body is
20 exposed to various forms of stress, including thermal, metabolic, hypoxic, oxidative, and
21 mechanical stress. These stressors activate biochemical messengers, which in turn activate
22 various signaling pathways that regulate gene expression and adaptive responses.
23 Historically, antioxidant supplements, nonsteroidal anti-inflammatory drugs, and
24 cryotherapy have been favored to attenuate or counteract exercise-induced oxidative stress
25 and inflammation. However, reactive oxygen species and inflammatory mediators are key
26 signaling molecules in muscle, and such strategies may mitigate adaptations to exercise.
27 Conversely, withholding dietary carbohydrate and restricting muscle blood flow during
28 exercise may augment adaptations to exercise. In this review article, we combine, integrate,
29 and apply knowledge about the fundamental mechanisms of exercise adaptation. We also
30 critically evaluate the rationale for using interventions that target these mechanisms under
31 the overarching concept of hormesis. There is currently insufficient evidence to establish
32 whether these treatments exert dose-dependent effects on muscle adaptation. However,
33 there appears to be some dissociation between the biochemical/molecular effects and
34 functional/performance outcomes of some of these treatments. Although several of these
35 treatments influence common kinases, transcription factors and proteins, it remains to be
36 determined if these interventions complement or negate each other, and whether such
37 effects are strong enough to influence adaptations to exercise.

38 Key words: adaptation, stress, preconditioning.

39

40 INTRODUCTION

41 Hormesis refers to ‘a process in which a low dose of a chemical agent or environmental
42 factor that is damaging at high doses induces an adaptive beneficial effect on the cell or
43 organism’ (127). The concept of hormesis first originated in the 16th century from the
44 musings of the Swiss physician and alchemist Paracelcus, who proposed that, “Solely the
45 dose determines that a thing is not a poison” (15). The term ‘hormesis’ itself was first coined
46 in 1943 by Southam and Ehrlich to explain their observation that a natural antibiotic in cedar
47 wood inhibited the growth of wood-decaying fungi but had the opposite effect at low doses
48 (204). Subsequently, the pioneering endocrinologist Hans Selye applied this notion to
49 understanding how biological systems respond to and tolerate environmental stress (194).

50 Hormesis encompasses the fundamental concepts of ‘conditioning’ and ‘adaptation’. The
51 concept of conditioning was first recognized following observations that repeated, brief
52 hypoxic exposure markedly reduced damage to the heart during subsequent myocardial
53 infarction (141). We now accept that exposure to an agent conditions the system to respond
54 in some manner (22). The concept of adaptation was originally recognized following
55 experiments demonstrating that constant exposure of *Escherichia coli* to mutagens allowed
56 each bacterium to handle mutagens more efficiently and to develop resistance to
57 mutagenesis (184). Conditioning and adaptation are closely related, are considered to be
58 synonymous, and are often used interchangeably. In essence, conditioning/adaptation
59 captures the notion that low levels of stress stimulate or upregulate existing cellular and
60 molecular pathways that improve the capacity of cells and organisms to withstand greater
61 stress (22).

62 The notion of hormesis underlies much of what we know about how exercise conditions
63 the body and induces long-term adaptation (32). However, hormesis was explicitly
64 introduced into the lexicon of exercise physiology only relatively recently (175). On a gross
65 population level, the dose–response nature of hormesis most likely explains why moderate
66 levels of physical activity reduce the risk of illness and mortality, whereas excessive physical
67 activity increases such risks (5, 103, 147).

68 During exercise, the body is exposed to various homeostatic perturbations, including
69 thermal, metabolic, hypoxic, oxidative, and mechanical stress. These perturbations
70 stimulate the release of biochemical messengers such as reactive oxygen and nitrogen
71 species (RONS), Ca^{2+} , growth factors, cytokines, and eicosanoids. These messengers then
72 activate signaling pathways including (but not limited to) various protein kinases,
73 phosphatases, and deacetylases, which in turn regulate the molecular machinery controlling
74 gene expression that elicits the appropriate adaptive responses (40). Through these
75 signaling pathways, acute production of RONS and inflammatory mediators can ultimately
76 promote adaptations in skeletal muscle such as mitochondrial biogenesis and
77 remodeling/hypertrophy (53, 124, 125, 169, 197). Conversely, prolonged production of
78 RONS and inflammatory mediators can activate proteolytic pathways, impede protein
79 synthesis, and overwhelm endogenous defense mechanisms, which cause adverse effects
80 such as muscle atrophy/weakness (37, 56, 62, 106, 203, 206). This dichotomy between the
81 acute and chronic effects of certain physiological stimuli is important to consider within the
82 context of hormesis in skeletal muscle.

83 Historically, the perception that exercise-induced oxidative stress and inflammation
84 cause muscle fatigue and damage has provoked widespread interest in countermeasures

such as antioxidant supplements, NSAIDs, and cryotherapy (31, 236). However, advances in our understanding of the role of RONS and inflammatory mediators in muscle adaptations to exercise have generated debate about whether these strategies are actually beneficial—at least in young healthy people (74, 162, 190). Antioxidant supplements and NSAIDs may help to preserve or enhance muscle adaptations to exercise in older individuals with impaired antioxidant defense systems or chronic low-grade inflammation (120, 228). By contrast, in young people these interventions can attenuate exercise-induced increases in insulin sensitivity (177) and muscle protein synthesis (229). The advantages or disadvantages of these interventions may therefore vary between different exercising populations. At the other end of the hormesis continuum, interest has also emerged in the potential benefits of applying stress to skeletal muscle before, during, or after exercise to stimulate greater adaptation. This stress can be applied by restricting carbohydrate intake, occluding local blood supply using low-intensity isometric or eccentric contractions (mechanical ‘preloading’), or passively heating muscle.

Considering the increasing attention on strategies to enhance exercise performance and assist recovery, it is timely to debate the scientific rationale for using interventions such as cryotherapy, antioxidant supplements, NSAIDs, mechanical preloading, dietary carbohydrate restriction, heat stress, and blood flow restriction to modulate adaptations to exercise. The purpose of this review is to combine, integrate, and apply knowledge about how these interventions influence skeletal muscle adaptations to exercise under the overarching concept of hormesis.

INTERVENTIONS THAT ENHANCE EXERCISE-INDUCED HORMESIS

108 *Restricting Dietary Carbohydrate Intake*

109 Modulating skeletal muscle glycogen content by restricting dietary carbohydrate intake
110 between exercise sessions is a relatively recent strategy to enhance exercise-induced
111 hormesis. Glycogen is an important substrate for oxidative phosphorylation in skeletal
112 muscle, and low muscle glycogen content is a key determinant of muscle fatigue (11, 94,
113 205). Accordingly, maximizing muscle glycogen content by carbohydrate loading before
114 exercise and delaying the rate at which glycogen content is depleted (by ingesting
115 carbohydrate during exercise) are common practices for athletes (21). Recent studies have
116 used various diet and/or exercise protocols to manipulate muscle glycogen content before
117 exercise sessions to determine whether changes in glycogen availability influence adaptive
118 responses [for review see (8, 69)]. There is growing evidence of beneficial effects on
119 metabolic and mitochondrial adaptations when exercising with low compared with normal
120 muscle glycogen content. This section briefly examines the putative mechanistic influence of
121 low muscle glycogen content and any potential for biphasic responses that support the
122 hormetic model of adaptation.

123 *Training with low muscle glycogen content promotes metabolic adaptation.* A primary
124 concept within the paradigm of nutrient–training interactions in skeletal muscle is that
125 substrate availability mediates the cellular response to contractile activity (32). However,
126 such a paradigm oversimplifies the complexity of how substrate availability modulates
127 adaptation. Hansen et al (65) first examined whether repeated bouts of exercise begun with
128 low muscle glycogen content induces greater metabolic stress and disruption to
129 homeostasis in skeletal muscle. They found that resting muscle glycogen content and citrate
130 synthase activity were higher in subjects who started half of their training sessions with low

glycogen versus those who always started training with normal glycogen. They concluded that this was because glycogen depletion caused by the first session dictated that the second session began with reduced muscle glycogen content. Although differences in the distribution of the training stimulus may have influenced these findings, there seems little doubt that the key factor promoting the adaptive response was training 'low'.

The metabolic flexibility of healthy skeletal muscle permits shifts in substrate oxidation based on the availability of carbohydrates and fats [for review see (205)]. Consequently, imposing the need for greater use of fat as a fuel likely explains much of the augmented adaptation to exercise with low initial muscle glycogen content. The demand for ATP supply during prolonged moderate- to high-intensity exercise is likely to also increase the magnitude of the adaptive signal under low-glycogen conditions. In this regard, the adenosine monophosphate activated protein kinase (AMPK) may be a focal point for regulating the cellular response to exercise with low initial muscle glycogen content, given its role as an energy sensor (66, 67). AMPK contains a glycogen-binding domain on one of its three subunits that causes it to colocalize with glycogen (66, 67, 128). AMPK also regulates the activity of several signaling pathways including those that promote glucose transport, fatty acid uptake, and mitochondrial biogenesis (66). The few studies that have quantified AMPK phosphorylation or activity after exercise begun with low- compared with normal/high-glycogen content have shown that the greater AMPK response in skeletal muscle following exercise is associated with lower preexercise glycogen content (242, 249).

Several other putative mediators of skeletal muscle adaptations to endurance exercise are enhanced after exercise with low initial muscle glycogen content. The phosphorylation status and mRNA abundance of important regulators of mitochondrial biogenesis (e.g.,

tumor suppressor p53 and peroxisome proliferator-activated receptor coactivator [PGC-1 α]) are more responsive to exercise with low compared with high initial muscle glycogen (9, 172). Similarly, mitochondrial enzyme activity increases after extended training periods during which exercise is repeatedly begun with low muscle glycogen content (65, 140, 250). Exercising with an initially low glycogen content also induces favorable metabolic responses, including greater oxidation of triacylglycerol and net uptake of glucose and fatty acids into skeletal muscle (75, 242, 250). Peroxisome proliferator-activated receptor δ expression increases in skeletal muscle after acute and chronic exercise (161), and likely plays an important function in alterations in muscle substrate metabolism following exercise training (17). Collectively, these findings suggest that manipulating carbohydrate availability before and/or during exercise stimulates several of the molecular and metabolic responses that promote adaptations to training.

Adverse responses to low glycogen content. Low glycogen availability limits its use for oxidative phosphorylation and may impair excitation–contraction coupling in muscle during exercise. Specifically, the reduction in Ca²⁺ release from the sarcoplasmic reticulum (SR) that accompanies muscle fatigue is associated with depletion of intramyofibrillar glycogen content (144, 256). In support of this *in situ* evidence, exercise studies have shown that depletion of muscle glycogen decreases Ca²⁺ release from the SR (50, 255). Importantly, SR Ca²⁺ release remains suppressed when carbohydrate intake is restricted in the early (4 h) postexercise recovery period. By contrast, resynthesis of muscle glycogen returns SR Ca²⁺ release rates to the preexercise levels (50). Together with the potential to promote shifts toward greater fat oxidation and inferior rates of carbohydrate oxidation, these responses could explain, at least in part, why acute exercise intensity is lower and endurance

177 performance following chronic training does not improve when using the ‘train low’
178 paradigm (75, 140, 250).

179 The increase in metabolic stress in skeletal muscle during exercise starting with low
180 glycogen content may also modulate protein turnover. In principle, higher AMPK activity
181 (resulting from low muscle glycogen content) could attenuate muscle protein synthesis by
182 inhibiting translation/elongation. Increased metabolic stress associated with low muscle
183 glycogen content may also exacerbate protein degradation (66, 72). Camera et al (23)
184 demonstrated that starting a bout of resistance exercise with low muscle glycogen content
185 neither promoted nor inhibited the myofibrillar protein synthesis. However, others have
186 reported that starting exercise with low muscle glycogen content increases the rates of
187 leucine oxidation and muscle protein degradation (13, 72). More research is needed to
188 determine the effects of training with low muscle glycogen content on protein turnover—
189 particularly during recovery between training sessions. Nevertheless, it is possible that
190 exercise starting with low compared with high muscle glycogen content may increase
191 muscle protein degradation.

192 Given the potential for conflicting beneficial and detrimental effects of training starting
193 with suboptimal glycogen content on skeletal muscle adaptations, a key question is: how
194 low should one go? If a biphasic response is dose dependent, one challenge is to titrate the
195 threshold for muscle glycogen content that might enhance the metabolic adaptations
196 without causing complications associated with fatigue or changes in the net protein balance
197 (Table 1). Perhaps the more pertinent question is not ‘how low’, but for ‘how long’ or ‘how
198 often’. Although acute restriction of dietary carbohydrate provides a positive stimulus for
199 metabolic adaptation, repeated depletion or long-term reduction in muscle glycogen

content may lead to overtraining (4). Therefore, the benefits of restricting carbohydrate during exercise or training with low initial muscle glycogen content must be balanced against the risk of fatigue.

Blood Flow-Restricted Exercise

In addition to nutritional interventions, it is also possible to enhance exercise-induced hormesis through physical interventions. One such example is applying a pressure cuff to the proximal regions of a limb during exercise. This practice first originated in Japan and was initially termed 'Kaatsu' training, which means 'adding pressure' (187). The first research published in English was a study by Shinohara et al (200), in which the combination of moderate resistance (40% of maximum voluntary contraction) and tourniquet ischemia resulted in a significant increase in strength (in contrast to no change in strength in the leg that exercised without ischemia). This training method is now more frequently referred to as 'blood flow-restricted exercise' (108). The basic physiological premise behind blood flow-restricted exercise is that it reduces blood flow and occludes the venous return from the limb (blood pooling). This combination of stimuli increases tissue hypoxia and the accumulation of metabolites, and thereby increases muscular stress during low-load resistance exercise (209-211). Blood flow-restricted exercise induces muscle hypertrophy and increases in muscle strength in the same range as traditional heavy-load strength training. Importantly, blood flow-restricted exercise induces effects that are absent (or minor) when low-load exercise is performed without blood flow restriction (108, 114).

221 Blood flow restriction results in several local and systemic responses that might
222 contribute to the enhanced hypertrophic stimulus when combined with low-load resistance
223 exercise [20–30% of 1 repetition maximum (RM)] (113, 240). In addition to metabolite
224 accumulation, the suggested mechanisms include increased recruitment of motor units
225 (rapid development of fatigue) (240), greater growth hormone secretion (215) and oxidative
226 stress (240), and muscle swelling (blood pooling) (110). Some of the mechanisms are
227 directly related, because metabolic accumulation causes rapid onset of fatigue (which
228 increases motor unit recruitment) and increases growth hormone secretion (215, 240).
229 Because it is difficult to separate these mechanisms, it remains unknown which of these
230 factors are most important. Nevertheless, combining blood flow restriction with low-load
231 resistance exercise increases the rate of muscle protein synthesis by activating similar
232 pathways to those activated after heavy-load strength training (e.g., mammalian target of
233 rapamycin [mTOR] signaling and MAPKs) (45, 47, 60, 239). Furthermore, low-load blood
234 flow-restricted exercise seems to induce a rapid and marked activation of satellite cells
235 (239). Interestingly, this satellite cell activation appears to exceed that which occurs after
236 traditional heavy-load strength training (145). Satellite cell activation induced by blood flow-
237 restricted exercise is accompanied by an increase in the number of myonuclei, which may
238 explain some of the muscle hypertrophy in response to blood flow-restricted exercise (18).
239 The 30–40% increase in cross-sectional area of both type I and II fibers after only seven
240 sessions of low-load, blood flow-restricted exercise supports the hypertrophic potential of
241 this method (145). Others have also reported rapid hypertrophy in response to high-
242 frequency (2×/day), low-load, blood flow-restricted exercise over 1–3 weeks (1, 3).

High-frequency, low-load blood flow-restricted exercise is generally a safe and effective training regimen because the low load induces less mechanical stress on muscle fibers than heavy-load strength training. In addition to the benefits described above, some studies also report no (or only minor) muscle damage and fast recovery after low-load, blood flow-restricted exercise (107, 112). However, the ischemia induced by blood flow restriction might cause some muscle damage and prolonged recovery if certain thresholds are passed. There are isolated reports of severe muscle damage resulting in rhabdomyolysis following blood flow-restricted exercise (82). Sarcolemmal and myofibrillar disruption and slow recovery of muscle function have also been reported after blood flow-restricted exercise in other studies (33, 241). These contrasting findings probably reflect differences in the training status of the study participants, degree of exhaustion, cuff pressure and size, and exercise intensity/volume.

Signs of damage, such as sarcolemmal disruption, high blood creatine kinase [CK] activity, and long-lasting fatigue, and rhabdomyolysis have been reported after the first session of low-load blood flow-restricted exercise (33, 82, 241), but rapid adaptation thereafter is likely. Performing a fixed number of repetitions per set (e.g., 15–15–15 or 30–15–15–15) causes little or no muscle damage (2, 111), but performing each set to failure causes more severe damage (33, 82, 241). The size of the cuff and the occlusion pressure can vary greatly. It can also be difficult to control arterial blood flow and venous return accurately (109). Collectively, these factors make it difficult to determine the optimal guidelines for blood flow restriction in combination with low-load resistance exercise.

Although the stress on the exercising muscle during low-load blood flow-restricted exercise is not well described, some interesting observations have been reported. In a

volume-matched protocol, blood flow-restricted exercise increased the acute expression of heat shock proteins (HSPs) in myofibrillar structures (33). Accumulation of small HSPs in myofibrillar structures was more abundant in type I fibers, indicating that low-load, blood flow-restricted exercise stresses type I fibers more than type II fibers, which contrasts with heavy-load strength training (43). This finding suggests that the combination of low-load resistance exercise and blood flow restriction preferentially stresses type I fibers. Provided that the stress remains within the optimal range, over the long term, such exercise also increases the hypertrophy of type I fibers. Importantly, in accordance with the hormesis theory, the dose is essential because excessive pressure and/or exercise volume/intensity may cause severe muscle damage, especially at the initiation of blood flow-restricted exercise.

In summary, applying a pressure cuff to restrict blood flow to an exercising limb—and thereby blocking venous return—increases the stress to the skeletal muscle during exercise. Blood flow restriction augments the effect of low-load resistance exercise on muscle hypertrophy. An important theme that arises from our evaluation is that blood flow restriction seems to shift muscular stress toward a more optimal range than that achieved with low-load exercise performed in isolation. However, the large variation in the application of blood flow restriction and exercise protocols makes it difficult to suggest an optimal protocol for low-load blood flow-restricted exercise at the present time. Acute blood flow restriction during exercise induces metabolic/hypoxic stress that ultimately leads to muscle hypertrophy. However, if used on a regular basis without sufficient recovery, blood flow-restricted exercise could induce a chronic cycle of muscle degradation and repair, which may impede rather than improve adaptations to training.

289

290 *Application of Heat to Muscle*

291 Applying heat to muscle is another physical intervention that may enhance exercise-
292 induced hormesis. Historically, heat has been used to treat severe muscle injuries (104),
293 although it may also improve recovery from less severe exercise-induced muscle damage.
294 The fundamental benefit of using heat in the management of muscle injuries involves an
295 increase in local blood flow (191, 245), which likely serves to improve the supply of oxygen
296 and nutrients to assist tissue repair (52). The alternative concept of using heat to
297 ‘precondition’ cells and tissues against other forms of stress was recognized around 20 years
298 ago. It was termed ‘cross-tolerance’ (248), and is a classic example of hormesis. It has
299 stimulated interest in the potential for heat preconditioning to protect myocardial tissue
300 against infarction (121) and skeletal muscle against atrophy (142). An increasing number of
301 studies have investigated the effects of heat application before or after various forms of
302 muscle injury on muscle regeneration and the associated mechanisms (Table 2).

303 *Heat preconditioning.* There is convincing evidence that heat stress assists recovery from
304 muscle injury. Application of heat (41°C) before *in vitro* muscle contraction augments
305 protein synthesis and expression of HSP72 in muscle cells (55, 247). In rats, heat
306 preconditioning 12–48 h before muscle injury increases muscle fiber cross-sectional area
307 and number of centrally nucleated fibers (96). This form of treatment also minimizes fiber
308 degeneration (199) and mitochondrial damage (48) after injury, and assists in maintaining
309 muscle mass during reloading after immobilization in rats (199).

Various mechanisms have been identified to explain these effects including: (i) an increase in phosphocreatine content, which is associated with less necrosis (48, 181); (ii) maintenance of reactive oxygen species-scavenging activity (199); (iii) increased expression of myosin heavy chain protein and HSPs (193, 223); and (iv) more Pax7⁺ satellite cells (96) in regenerating muscle. Heat preconditioning also reduces oxidative damage to muscle protein (193) and infiltration of mononuclear inflammatory cells (96, 199, 223) after muscle injury in rats. In addition to these studies on muscle injury, heat preconditioning increases the activity of PGC-1 α and AMPK in C2C12 myotubes (84) and prevents muscle atrophy in response to immobilization (192) and hindlimb unloading in rats (142).

Research on the effects of heat preconditioning on recovery from exercise-induced muscle damage in humans has produced more variable findings. Some work indicates that heat stress before eccentric exercise can reduce muscle fatigue (77), promote faster recovery of strength and range of motion, and alleviate muscle soreness (149, 183). Heat preconditioning also increases the activation of Akt, mTOR, ribosomal protein S6, and eukaryotic translation initiation factor 4E-binding protein 1 (EIF4E-BP1) after resistance exercise (90). In contrast with these studies, others have reported no benefits of heat preconditioning on the recovery of strength, range of motion, edema, or soreness after eccentric exercise (86, 151).

Heat stress after muscle injury/exercise. Various animal studies have reported that applying heat after muscle injury increases muscle fiber cross-sectional area and number of centrally nucleated fibers (68, 71, 96, 154, 216). Consistent with the effects of heat preconditioning, these benefits of therapeutic heat treatment are conferred by upregulation of HSPs in muscle (71, 154). Heat application after muscle injury in rats also induces more rapid

333 macrophage infiltration (216); expression of IGF-1 (216), MyoD, and myogenin (68),
334 calcineurin (154); and activity of Pax7⁺, MyoD⁺, and M-cadherin⁺ satellite cells (96, 154, 216).
335 Conversely, applying heat to muscle following injury reduces myeloperoxidase activity,
336 production of RONS, lipid peroxidation, and fibrosis in rats (25, 71, 216).

337 Relatively little is known about how applying heat to muscle after exercise influences
338 acute recovery of muscle function. One study reported that, compared with passive
339 recovery, hot water immersion (38°C for 14 min) after eccentric exercise improved the
340 recovery of strength, but not that of muscle power, swelling, or soreness (231). The same
341 group reported that hot water immersion did not help to maintain sprint or time trial
342 performance over 5 days of high-intensity cycling (230).

343 No studies have investigated the effects of regular heat application on chronic muscle
344 adaptations to training. However, evidence from a recent study on rats suggests some
345 potential benefits of heat to enhance training adaptations. In this study, rats that were
346 placed in a heat chamber at 41°C for 30 min immediately after treadmill running showed
347 greater chronic increases in the activity of citrate synthase and 3-hydroxyacyl CoA
348 dehydrogenase, and mitochondrial protein content in skeletal muscle after 3 weeks of
349 training (5 days/week) (217).

350 The transcription factor heat shock factor-1 (HSF-1) and its downstream effectors, HSPs,
351 are most likely central to the benefits of heat stress for healing of muscle injuries, as
352 demonstrated in animal studies outlined below. HSF-1 and HSPs may assist muscle
353 regeneration by protecting muscle cells against oxidative damage, apoptosis, and ATP
354 depletion (16, 87-89, 118). HSPs may also promote repair of muscle tissue by activating the
355 signaling pathways involved in protein synthesis (e.g., Akt, p70S6 kinase, and ERK) (61) and

by regulating the activity of enzymes and transcription factors that can cause degeneration and/or atrophy of muscle fibers (38, 49, 105, 195). Importantly, without HSF-1 and HSP70, macrophage infiltration is delayed, and the expression of proinflammatory cytokines is dysregulated in regenerating muscle tissue (98, 148, 196). Heat stress may also increase muscle hypertrophy independently of HSPs by stimulating the expression of IGF-1, myogenin, and Pax7 (166). Increased expression of IGF-1 in response to heat stress likely complements the effects HSPs by orchestrating more efficient resolution of inflammation following muscle injury (160).

This review is the first summary and critical evaluation of the effects of applying heat to muscle with the goal of promoting repair and growth of muscle. Acute heat stress increases the activities of HSPs, satellite cells, PGC-1 α , and AMPK, whereas it reduces oxidative damage in muscle after exercise/injury. Over the long term, these responses may augment training adaptations. Although the application of heat stress before or after muscle injury has shown promising results in muscle cell culture and animal studies, more work is required to establish whether these same benefits occur in humans.

Mechanical Preloading

A single bout of eccentric muscle contractions confers protection against subsequent bouts of muscle-damaging exercise. This response is referred to as the 'repeated-bout effect', and may last between 6 and 9 months (150). The repeated bout effect can also occur in the non-exercising contralateral limb, although the effect in the contralateral limb is smaller than that in the ipsilateral limb (73).

Recent interest has focused on trying to determine the minimum stimulus required to elicit protection against muscle damage, which is typically characterized by prolonged decreases (>1 d) in muscle function and delayed-onset muscle soreness (DOMS). Herein, we refer to this approach to strength training and conditioning as ‘mechanical preloading’. Although this is a relatively new concept, it is a classic example of exercise-induced hormesis, whereby mild mechanical preloading of skeletal muscle induces positive adaptations. The first evidence for the benefits of mechanical preloading came from a study demonstrating that low-intensity isometric contractions (performed at 10% of maximal voluntary contraction strength) improved the recovery of strength by 50–60% and reduced peak muscle soreness by 30% after subsequent eccentric exercise performed 2 days later (101). These protective effects of mechanical pre-loading seem to last between 1 and 2 weeks (26).

Mode and intensity of contraction. The preloading effect does not appear to be specific to the type of muscle contraction. Preloading with as few as two maximum voluntary isometric contractions at a long muscle length (20° flexion) is sufficient to attenuate the loss of strength and range of motion, DOMS, and swelling after eccentric exercise performed 2 days later (27). As evidence of a dose response, 10 maximal voluntary isometric contractions at the same muscle length conferred even greater protective effects (27). The protective effect conferred by two maximal isometric contractions appears to last only a maximum of 1 week (28). Compared with low-intensity eccentric contractions (10% maximum strength), maximal isometric contractions performed at 20° flexion confer a greater degree of protection against subsequent muscle damage (30). However, the protective effect of maximal isometric contractions is less than that resulting from maximal eccentric

contractions (30). Four bouts of moderate-intensity eccentric exercise comprising eccentric contractions at 40% of maximal voluntary isometric contraction, performed every 2 weeks, confers a similar protective effect to one bout of maximal eccentric exercise (29). This finding suggests that repeating submaximal eccentric exercise provides the same protection as one bout of maximal eccentric exercise against the subsequent maximal eccentric exercise. It remains to be determined whether regular lighter intensity eccentric contractions (e.g., 10%) or maximal isometric contractions at a long muscle length increase long-term muscle adaptations.

Integration of the findings of the small number of studies in this area shows that a few eccentric contractions at low intensity or a few maximal isometric contractions at long muscle length confer significant protection against subsequent muscle damage. In addition to contracting muscles, this effect most likely also occurs in non-exercising muscles of the contralateral limb. The mechanisms underpinning the effects of mechanical preloading on muscle adaptation are currently unknown. Adaptation to maximal eccentric contractions has been attributed to various factors, including neural changes (e.g., increased motor unit recruitment/synchronization), remodeling of connective tissue, removal of weak fibers, and longitudinal addition of sarcomeres (131). Light-intensity eccentric contractions and isometric contractions do not cause any loss of strength or range of motion, muscle swelling, or DOMS (27, 101). Without causing frank muscle damage, these types of contractions may precondition skeletal muscle through other mechanisms. Such mechanisms could include physical changes to the fascia and endomysium or metabolic alterations in ATP availability, intracellular $[Ca^{2+}]$, mitochondrial Ca^{2+} uptake, RONS signaling,

or proteolytic activity. Further research is warranted to examine these putative mechanisms in greater detail.

Because acute muscle damage resulting from mechanical preloading is minimal, it seems unlikely that long-term use of this form of preconditioning will increase the risk of maladaptation to training. However, the protective effect of mechanical preloading may diminish if it is used repeatedly because muscle probably adapts to such mechanical stimulation. Consistent with this premise, any benefits of mechanical preloading are probably relatively minor for resistance-trained individuals who regularly perform submaximal eccentric contractions and maximal isometric contractions in their training routines. Future studies in this area could investigate whether skeletal muscle remodeling/hypertrophy is still induced effectively if no muscle damage is induced throughout training.

INTERVENTIONS THAT DAMPEN EXERCISE-INDUCED HORMESIS

Antioxidant Supplementation

The notion of hormesis has been studied extensively in the context of oxidative stress and its opposing roles in skeletal muscle pathologies. It has also been examined as a potential stimulus for redox adaptations in skeletal muscle following endurance training. For the purposes of this review, the term ‘oxidative stress’ is defined as an imbalance between oxidants and antioxidants in favor of the oxidants, leading to a disruption of redox signaling and control and/or molecular damage (171). Davies et al (34) were the first to report that submaximal exercise to exhaustion increased the production of free radicals in rodent

skeletal muscle. Other more recent studies have also shown that exhaustive endurance exercise increases oxidative stress in rat skeletal muscle (10, 91, 235). Although these studies provide vital proof of principle, understanding precisely how RONS regulate skeletal muscle adaptations to endurance training is difficult—mainly because few training programs regularly push individuals to exhaustion. Nevertheless, moderate- to high-intensity endurance exercise (70–85% of maximal oxygen uptake) is sufficient to increase oxidative stress in rat skeletal muscle, as measured by changes in GSSG levels (237, 238, 254). Moderate-intensity endurance cycling exercise is also sufficient to increase lipid peroxidation, as measured by F₂-isoprostane content in skeletal muscle of humans (92). Bailey et al (7) provided the first direct evidence in humans that exercise in the form of maximal, single-leg knee extension increases intramuscular free radical accumulation.

Oxidative stress and mitochondrial biogenesis in skeletal muscle. Redox-sensitive kinases activated during muscle contraction include AMPK, activating transcription factor-2 (ATF-2), NFκB, and the MAP kinases p38 MAPK, JNK, and ERK (also called p44/42 MAPK) (53, 79, 185, 238). These kinases are all implicated in the regulation of mitochondrial biogenesis (83, 243)—at least partly through the transcriptional coactivator PGC-1α, which is a key regulator of mitochondrial biogenesis (173, 244). Although RONS were first proposed to regulate exercise-induced mitochondrial biogenesis over 30 years ago (34), it was Silveira et al who first published clear evidence linking RONS with the regulation of contraction-induced mitochondrial biogenesis in rat muscle cells (201). Importantly, this group demonstrated that antioxidants attenuated the increase in RONS production and PGC-1α mRNA expression (201). Hood et al (79) have since provided more direct evidence for the role of RONS (and antioxidants) in regulating the expression of AMPK and PGC-1α in skeletal

muscle cells. Other proteins such as upstream stimulatory factor 1 also play an important role in regulating PGC-1 α activity in skeletal muscle (80).

Antioxidants and mitochondrial biogenesis. Research on the effects of antioxidants on mitochondrial biogenesis has used vitamins C and E (alone or in combination), coenzyme Q10, *N*-acetylcysteine, β -carotene and α -lipoic acid in rats (54, 70, 208, 234) and humans (157, 163, 177, 251). Because of the large number of individual antioxidant supplements, a comprehensive examination of each antioxidant is beyond the scope of the current review (for review, see (120). This review is limited to evaluation of hormesis specifically in relation to vitamins C and E because they are two of the most common antioxidant supplements used alone or in combination by the general population (180) and in research (54, 70, 157, 177, 208, 234, 251). Given the role of RONS in stimulating mitochondrial biogenesis in skeletal muscle (79, 201), many studies have investigated whether antioxidant supplements prevent adaptations to endurance training. Some training studies have found that vitamin C and/or vitamin E attenuates markers of mitochondrial biogenesis in muscle after training in rats (54, 234) and humans (157, 177). By contrast, other studies have found no significant effects of antioxidant supplements on markers of mitochondrial biogenesis (70, 208, 251, 253).

Despite this evidence for a reduction in cellular adaptations to endurance training with antioxidants (54, 157, 177, 234), no research has reported any change in maximum oxygen uptake or exercise performance—at least in humans (157, 178, 251). Animal studies have demonstrated that vitamin C supplementation reduces the improvements in exercise performance after 6 weeks of exercise training (54, 126). Differences in the metabolism of

490 vitamin C in skeletal muscle between humans and rats may partially account for these
491 differences.

492 Despite strong evidence that endurance exercise increases oxidative stress in human
493 skeletal muscle (7, 92, 254), it remains uncertain whether vitamin C and/or E
494 supplementation inhibits oxidative stress in human skeletal muscle during exercise. One
495 reason for this uncertainty is the lack of suitable markers of RONS production and oxidative
496 stress in skeletal muscle during exercise. Some studies have used plasma or blood to assess
497 oxidative stress (70, 157). However, this is problematic because the degree of systemic
498 oxidative stress in plasma/blood may not reflect the extent of local oxidative stress in
499 skeletal muscle (235). Furthermore, other markers of oxidative stress (e.g., thiobarbituric
500 acid reactive substances (TBARS) or malondialdehyde) may not be specific or sensitive to
501 antioxidant supplementation (182, 252).

502 In addition to discrepancies between the effects of antioxidants in animals compared
503 with humans, there is also some disparity between the acute and chronic effects of
504 antioxidants. For example, several acute exercise studies show that inhibiting RONS derived
505 from xanthine oxidase with the xanthine oxidase inhibitor, allopurinol, inhibits the exercise-
506 induced phosphorylation of redox-sensitive kinases such as p38 MAPK and ERK, which
507 regulate mitochondrial biogenesis in rats (53, 91, 238). However, long-term treatment with
508 allopurinol does not prevent the increases in skeletal muscle mitochondrial proteins or
509 antioxidant enzymes following endurance training in rats (238). One possible reason for this
510 disparity is that stimuli other than RONS, such as cytosolic Ca^{2+} (130, 155), AMP (130), and
511 possibly NAD (51) also regulate mitochondrial biogenesis in skeletal muscle. Thus, although
512 antioxidant supplements can inhibit RONS production in skeletal muscle, this may not

513 always attenuate mitochondrial biogenesis probably because of redundancies within these
514 pathways.

515 *Antioxidants and skeletal muscle hypertrophy.* There is substantial evidence linking oxidative
516 stress with muscle atrophy [for review see (170)]. Emerging evidence also implicates
517 oxidative stress in the regulation of skeletal muscle hypertrophy. A high daily oral dose of
518 vitamin C attenuates skeletal muscle hypertrophy and oxidative stress normally observed
519 following mechanical overload of the plantaris (119). Recent findings in rodents
520 demonstrate that the highly reactive oxidant, peroxynitrite regulates skeletal muscle
521 hypertrophy induced by overload (81). Peroxynitrite appears to operate by stimulating the
522 release of intracellular Ca^{2+} , which then activates mTOR to increase protein synthesis (119).

523 The few human studies to investigate the adaptations to resistance training combined
524 with antioxidant supplementation have reported variable findings. Two studies showed no
525 effect of vitamin C and E supplementation on improvements in skeletal muscle strength or
526 performance (14, 220). However, these studies used resistance training protocols that did
527 not induce skeletal muscle hypertrophy (14) or did not measure changes in lean muscle
528 mass (220). Paulsen et al (159) recently found that supplementation with vitamins C and E
529 attenuated the activities of several kinases involved in hypertrophy signaling, such as p70S6
530 kinase and the redox-sensitive kinases p38 MAPK and ERK 1/2 in skeletal muscle after 10
531 weeks of resistance training. In addition, supplementation attenuated bicep curl strength
532 following 10 weeks of training. By contrast, supplementation did not alter protein synthesis
533 or muscle hypertrophy following training (159). Thus, some evidence supports blunting of
534 the cell signaling pathways with antioxidant supplementation following resistance exercise,
535 although the effects on functional outcomes remain equivocal. More studies are required to

examine whether RONS regulate hypertrophy following resistance training in human skeletal muscle and whether antioxidant supplementation influences these adaptations to resistance exercise.

In summary, oxidative stress plays an important role in regulating the mitochondrial content and perhaps contractile protein content of skeletal muscle. Some evidence shows that supplementation with vitamins C and E can block acute increases in signaling pathways that control mitochondrial biogenesis and hypertrophy. However, these acute responses do not consistently translate to less mitochondrial biogenesis or muscle hypertrophy following chronic exercise training because of the apparent redundancy in skeletal muscle. That is, exercise training (either endurance or resistance) may induce mitochondrial biogenesis and hypertrophy despite elevated concentrations of RONS-scavenging antioxidants. The weight of current evidence suggests that vitamin C and E supplementation may dampen exercise-induced hormesis—at least at the cellular level. However, it remains uncertain whether these responses influence exercise performance in the long term. Importantly, antioxidant compounds have widely divergent properties, and this discussion of a specific class of agents does not rule out the effects of other components on RONS activity/regulation, nor a role for RONS in exercise-induced adaptation. The requirement for and efficacy of antioxidant supplements may vary with age and health status. There are conflicting and unresolved issues surrounding the influence of antioxidant supplementation on adaptations to training that require further investigation.

NSAIDs

Similar to antioxidants, NSAIDs represent another pharmacological intervention that may attenuate exercise-induced hormesis. NSAIDs are inhibitors of the cyclooxygenase (COX) pathway that converts free arachidonic acid to PGD₂, PGE₂, PGF_{2α}, PGI₂, and thromboxane A₂ (42, 232). PGs are autocrine/paracrine lipid mediators that propagate the inflammatory response to tissue injury by increasing blood flow, vascular permeability, and leukocyte chemotaxis (35). COX has two major isoforms. COX-1 is constitutively expressed, and COX-2 expression is generally low but is highly inducible in response to injurious stimuli (57, 139). Classical NSAIDs inhibit both COX-1 and COX-2 to varying degrees (36, 202). Undesirable side effects associated with disruption of homeostatic COX-1 activity have led to the development COX-2-specific inhibitors (coxibs) for treating pain and inflammation. During postexercise recovery, the activities of COX-1 and COX-2 (24) and concentrations of PGs (20, 93, 225, 227) increase transiently in skeletal muscle. Plasma PG concentrations also increase after exercise (39, 123, 218). These responses point to important roles for the COX/PG pathway in exercise adaptation. On the other hand, chronically elevated PG concentrations are associated with—and may contribute directly to—muscle wasting in states of chronic inflammation (97).

Effect of NSAIDs on acute muscle responses to exercise. Classical NSAIDs (e.g., ibuprofen and indomethacin) administered at over-the-counter doses effectively block the acute exercise-induced increase in PG concentration in muscle (20, 135, 227) and plasma (123). Although not considered a classical NSAID, acetaminophen also appears to inhibit COX activity in muscle (227). Many studies have investigated the effect of NSAIDs on symptoms of exercise-induced muscle damage, although the literature on the efficacy of NSAIDs for reducing muscle soreness and/or improving exercise recovery is contradictory. Given that NSAIDs are

anti-inflammatory, it is surprising that studies to date have failed to observe any effect of NSAIDs on systemic (95, 167, 222) or intramuscular (158) leukocyte responses to exercise stress. Paradoxically, short-term NSAID treatment appears to increase plasma cytokine concentrations (e.g., IL-6 and monocyte chemotactic protein-1) (41, 59, 138, 146) and muscle COX-2 gene expression (19, 138) after exercise.

Together with a lack of a clear benefit of NSAIDs in reducing exercise-induced pain and/or the acute inflammatory response in humans, various studies have shown potential negative effects of NSAIDs in muscle after exercise. Oral ingestion of the nonselective NSAIDs ibuprofen or acetaminophen blunts the increase in muscle protein synthesis during postexercise recovery in young men (229). However, this effect was not replicated in a study of patients with knee osteoarthritis who received ibuprofen (165). Another nonselective NSAID (indomethacin) blocked the muscle satellite cell response to a 36 km run (117) and maximal eccentric exercise (137) but did not alter muscle protein synthesis (138). Studies have shown that COX-2-selective inhibitors do not influence muscle protein synthesis (19) or satellite cell responses to exercise (158), suggesting that COX-1 rather than COX-2 may be the primary isoform involved in human muscle responses to exercise.

The underlying mechanisms by which NSAIDs influence muscle adaptive responses to exercise remain unclear, but several recent studies have provided useful insights. Impaired satellite cell proliferation following maximal eccentric exercise with local indomethacin infusion (135) did not alter the expression of growth factors and extracellular matrix-related genes (138) or HSP (136) in muscle. Oral ibuprofen treatment blocked the normal increase in serum PG concentration during early postexercise recovery (0–3 h) (123), and suppressed phosphorylation of components of the ERK and mTOR signaling pathways in muscle (122).

These data provide the first evidence that PGs contribute to contraction-induced signaling in human muscle and provide mechanistic support for a potentially detrimental effect of oral nonselective NSAIDs (122, 125). Interestingly, mass spectrometry profiling of serum samples collected throughout exercise recovery revealed suppression of both early proinflammatory and later anti-inflammatory/proresolving lipid mediator circuits in subjects receiving ibuprofen (123). Thus, NSAIDs may interfere with exercise recovery indirectly by delaying or preventing timely resolution of the inflammatory response (123, 233).

Chronic effects of NSAIDs on muscle exercise adaptation. Although nonselective NSAIDs may attenuate acute responses to exercise in humans (122, 123, 137, 138, 227, 229), it remains unclear whether these responses influence long-term adaptations to exercise. Oral ibuprofen treatment (400 mg/day) did not influence muscle hypertrophy or strength following 6 weeks of resistance training of the elbow flexors in young healthy men (99). However, this dose of ibuprofen was only one-third that used in acute exercise studies (122, 123, 227, 229). By contrast, animal studies clearly show a deleterious effect of NSAID treatment on long-term muscle regeneration and hypertrophy, and specifically implicate the COX-2 isoform in this response (100, 124, 152, 198).

In older adult subjects, gains in skeletal muscle size and strength following 12 weeks of resistance training were greater in response to treatment with ibuprofen (1,200 mg/day) or acetaminophen (4 g/day) compared with a placebo treatment (224). Another study also revealed that ibuprofen augmented training-induced gains in muscle strength in elderly subjects but did not influence muscle mass and tended to reduce satellite cell numbers in muscle (164). By contrast, a lower dose of acetaminophen (1,000 mg/day) did not alter fat-free-mass or muscle strength in older men after a period of resistance exercise training (85).

One mechanism through which NSAIDs may exert positive effects on muscle involves a reduction in chronic low-grade inflammation that occurs with aging, thereby blocking the pathway to muscle atrophy. NSAID treatment counteracts skeletal muscle wasting in animal models of chronic inflammatory disease including cancer cachexia (129, 202, 207), arthritis (56), and aging (176). Consistent with this hypothesis, older adults who received ibuprofen throughout 12 weeks of resistance training showed a chronic reduction in the expression of cytokine genes (e.g., IL-6, IL-10) and muscle ring finger 1 (MuRF-1) (226).

In summary, the COX/PG pathway appears to play an important role in acute exercise recovery, and NSAIDs inhibit the seemingly beneficial acute muscle adaptive responses to exercise (e.g., satellite cell proliferation and muscle protein synthesis). On the other hand, chronic activation of the COX/PG pathway may exert negative effects on muscle mass, and NSAID treatment may provide an effect countermeasure against such effects. In this review, we have highlighted an apparent discrepancy between the opposing effects of NSAIDs in different settings (e.g., acute versus chronic, young versus old subjects). The balance between PG species with differing bioactivity (e.g. $\text{PGF}_{2\alpha}$ versus PGE_2) (228) or differences in the underlying nature of the inflammatory response (acute self-resolving versus chronic nonresolving) (97, 122) may be important factors that influence the pharmacological actions of NSAIDs.

Cryotherapy

Cryotherapy in the form of ice massage and application of crushed ice has long been a common treatment for soft tissue injuries (132). More recently, other forms of cryotherapy

such as cold water/ice baths and brief exposure to extreme cold air (–20 to –110°C) in custom-made cryotherapy chambers have gained popularity as strategies to recover from exercise. Traditionally, the physiological basis for using cryotherapy has been to relieve pain, reduce tissue metabolism, and modify vascular responses to minimize edema (213). Acute responses to primary muscle injury (e.g., necrosis and inflammation) can result in ‘secondary injury’ to healthy cells not damaged through the initial trauma (134). By reducing the metabolic rate of tissues within and around the injury site, cryotherapy may protect the healthy bystander cells from the ischemic environment in the immediate period after injury, thereby reducing the risk of secondary cell injury or death (12). Some evidence from animal studies support this notion (133, 134, 156, 186). However, the effects of cryotherapy on muscle inflammation in humans are currently unknown.

Effects of cryotherapy on inflammation and oxidative stress. Studies have focused on how icing influences inflammation and oxidative stress in muscle following injury (Table 3). Superfusing rats with cold saline (3–8°C) for 10 min to 6 h after muscle contusion injury significantly reduced leukocyte rolling and adhesion to venules within damaged muscle for up to 1 day after injury (102, 188, 189). These effects may be mediated by downregulation of adhesion molecules on the surface of vessels and leukocytes in response to hypothermia (63, 78). Immunohistochemical analysis of muscle tissue revealed that this cryotherapy treatment decreased the number of neutrophils in muscle 1 day after injury (188, 189). In support of these findings, others have observed that icing after muscle strain injury in rats substantially reduced neutrophil activation in muscle, as indicated by lower myeloperoxidase activity 1 day after injury (25). Icing also restricted the production of RONS and lipid peroxidation at 1, 5, 10 and 15 days after injury in rats (25). Icing preserves the

activity of $\text{Na}^+\text{-K}^+\text{-ATPase}$ and $\text{Ca}^{2+}\text{-ATPase}$ enzymes and mitochondrial membrane permeability, and it reduces mitochondrial swelling in muscle 1 day after contusion injury in rats (174). Because none of these studies assessed muscle regeneration in the weeks following injury, it is difficult to establish whether restricting neutrophil invasion and activation through cryotherapy results in better healing of muscle injuries. In principle, a decrease in neutrophil infiltration into muscle as a result of icing is potentially beneficial because activated neutrophils can damage skeletal muscle fibers (143, 168).

Effects of cryotherapy on muscle regeneration. Other studies in rats have shown that icing causes greater fibrosis and impairs muscle regeneration after muscle contusion and crush injuries. These effects are evident as early as 2 days after injury (76) and persist for up to 4 weeks (214). The potential mechanisms responsible for these effects include delayed macrophage infiltration and mRNA expression of transforming growth factor- β 1 and IGF-1 in muscle, together with a delay in (or absence of) satellite cell activation (76, 214). Impaired muscle regeneration in response to icing may be attributed to the following sequence of events. By restricting neutrophil infiltration, icing may slow the rate of phagocytosis of necrotic muscle tissue in the first few hours after injury (219). Persistent necrosis may then delay the entry of macrophages into muscle tissue in the first few days after injury (58). Finally, by delaying macrophage infiltration, icing may reduce the capacity of these cells to (a) produce essential growth factors and chemotactic agents (64, 115, 116, 212), and (b) stimulate satellite cells to proliferate and differentiate (6, 221). The limited evidence that is currently available therefore suggests that cryotherapy is detrimental for muscle regeneration following injury.

Effects of cryotherapy on training adaptations. In addition to this research on acute muscle injury, a smaller body of research has investigated the effects of regular cryotherapy on muscle adaptations to exercise training. An early study demonstrated that, in rats regularly immersed in cold water (4°C) for 5 min after exercise bouts, greater ultrastructural damage to myofibrils was evident after 5 weeks of exhaustive running and 7 weeks of moderate running (46). Fu et al proposed that, by masking pain, cold water immersion allowed the rats to exercise at higher intensities the next day, which unexpectedly resulted in greater muscle damage (46). Subsequently, several human studies have also reported that regular cold water immersion after exercise attenuates muscle adaptations to training (44, 153, 179, 246). The mechanisms by which regular cold water immersion dampened training adaptations in these studies are unknown. Hypothetically, a decrease in muscle blood flow in response to cold water immersion might reduce angiogenesis and protein synthesis in muscle during recovery from exercise. In turn, these responses may result in smaller gains in muscular endurance and strength.

This review is the first critical evaluation of the short- and long-term effects of various forms of cryotherapy on cellular responses in skeletal muscle. We have also outlined in detail the putative mechanisms by which cryotherapy influences muscle repair and growth. When applied acutely after exercise or muscle injury, cryotherapy may help to reduce muscle soreness and minimize secondary tissue damage. However, by attenuating some key inflammatory reactions (e.g., macrophage infiltration) in skeletal muscle, cryotherapy may also block the production and release of important growth factors and the activity of satellite cells, which are important mediators of muscle repair and adaptation. Therefore,

although cryotherapy offers some short-term benefits, these are possibly outweighed by long-term detrimental effects.

PERSPECTIVES AND FUTURE DIRECTIONS

This is the first commentary to combine, summarize, and evaluate the efficacy of various strategies to modulate exercise-induced hormesis. Some of these strategies (e.g., antioxidant supplementation, treatment with NSAIDs, restriction of dietary carbohydrate intake) have been the subject of scientific scrutiny and debate. By contrast, other strategies such as cryotherapy, blood flow restriction, heat stress, and mechanical preloading have received less critical attention. In this review, we have detailed the conceptual frameworks for the use of such strategies, have integrated these details with the current knowledge about the basic biochemical and molecular machinery that regulate muscle adaptations to exercise, and have applied this information to assess the advantages and disadvantages of each strategy for modulating exercise-induced hormesis.

Table 4 summarizes the mechanisms of action of treatments that modulate exercise-induced hormesis and describes some of the short- and long-term outcomes of these treatments. A key finding from this review is that there appears to be some dissociation between the biochemical/molecular effects and functional/performance outcomes of some of these treatments (e.g., antioxidants, NSAIDs, restriction of dietary carbohydrate). Conceivably, other signaling pathways that are less responsive to these treatments (or not yet defined) may operate independently in the regulation of training adaptations. This redundancy may promote fine-tuning of adaptive responses to exercise training (40). Few of

the interventions described in this review have been adequately tested to determine if or how they exert dose-dependent effects on muscle adaptation. If such dose-dependent effects do occur, they are likely to be subject to highly complex regulatory mechanisms.

A common feature of hormesis is that exposure to one type of hormetic agent can protect cells/organisms against more types of stress (127). This concept of 'cross tolerance' may be applied to some of the interventions that we have discussed. Several of the interventions influence common kinases, transcription factors, and proteins (see Table 4). For example, AMPK, p38 MAPK, PGC-1 α , and HSP expression increases in response to heat stress, carbohydrate restriction, and blood flow restriction, whereas the expression of most of these factors decreases following antioxidant supplementation. Similarly, macrophage infiltration, IGF-1, and Pax7 expression increases in response to heat stress, whereas these factors are either blocked or activated more slowly after cryotherapy. It remains to be determined whether these interventions complement or negate each other and whether such effects are strong enough to alter terminal adaptive processes such as mitochondrial biogenesis, substrate metabolism, or muscle repair/growth.

Several important questions have emerged from this review that warrant further investigation. A primary issue relates to the threshold (i.e., dose, period of exposure) that defines whether oxidative stress and inflammation are beneficial for or harmful to muscle adaptations to exercise. This threshold would be difficult to titrate because it most likely depends on the basal state of oxidative stress and inflammation at the start of exercise. In turn, this basal state may depend on periodization of training and recovery, together with age, health status, and diet. In addition, it is unclear whether undertaking different strategies simultaneously enhances or attenuates exercise-induced hormesis and which

combination of strategies might offer complementary or additive benefits. As highlighted in our review, some interventions such as NSAIDs and antioxidants exert different effects in young compared with older individuals and in trained compared with untrained individuals. Finally, the efficacy of a given intervention may depend on the capacity to 'periodize' such interventions during different phases of a training program. For example, during training to promote muscle hypertrophy and strength, interventions such as cryotherapy and the use of NSAIDs may dampen rather than enhance adaptation. However, during periods of regular competition when recovery is a priority, these strategies may be appropriate to alleviate muscle soreness and restrict secondary tissue injury.

In conclusion, exercise-induced adaptations in skeletal muscle are regulated through interactions between various mechanical, metabolic, and physiological stressors and complex cellular machinery. Undoubtedly, a large body of work is still required to provide greater clarity on the appropriate uses and applications of strategies to modify skeletal muscle phenotypes. Exercise-induced hormesis is an intriguing notion that awaits further exploration. To adapt a phrase from a well-known bard, to intervene or not intervene: that remains the question.

777

778 **References**

- 779 1. **Abe T, Kawamoto K, Yasuda T, Kearns CF, Midorikawa T, Sato Y.** Eight days KAATSU–
780 resistance training improved sprint but not jump performance in collegiate male track and
781 field athletes. *Int J Kaatsu Training Res* 1: 19–23, 2005.
- 782 2. **Abe T, Yasuda T, Midorikawa T, Sato Y, Kearns C, Inoue K, Koizumi K, Ishii N.** Strength
783 training with low loads in combination with vascular occlusion has been proposed as an
784 alternative to heavy resistance exercise. *Int J Kaatsu Training Res* 1: 6–12, 2005.
- 785 3. **Abe T, Yasuda T, Midorikawa T, Sato Y, Kearns CF, Inoue K, Koizumi K, Ishii N.** Skeletal
786 muscle size and circulating IGF–1 are increased after two weeks of twice daily kaatsu
787 resistance training. *Int J Kaatsu Training Res* 1: 7–14, 2005.
- 788 4. **Achten J, Halson SL, Moseley L, Rayson MP, Casey A, Jeukendrup AE.** Higher dietary
789 carbohydrate content during intensified running training results in better maintenance of
790 performance and mood state. *J Appl Physiol* 96: 1331–1340, 2004.
- 791 5. **Andersen K, Farahmand B, Ahlbom A, Held C, Ljunghall S, Michaelsson K, Sundstrom J.**
792 Risk of arrhythmias in 52 755 long-distance cross-country skiers: a cohort study. *Eur Heart J*
793 34: 3624–3631, 2013.
- 794 6. **Arnold L, Henry A, Poron F, Baba-Amer Y, van Rooijen N, Plonquet A, Gherardi RK,**
795 **Chazaud B.** Inflammatory monocytes recruited after skeletal muscle injury switch into
796 antiinflammatory macrophages to support myogenesis. *J Exp Med* 204: 1057–1069, 2007.
- 797 7. **Bailey DM, Lawrenson L, McEneny J, Young IS, James PE, Jackson SK, Henry RR,**
798 **Mathieu-Costello O, McCord JM, Richardson RS.** Electron paramagnetic spectroscopic
799 evidence of exercise-induced free radical accumulation in human skeletal muscle. *Free Radic*
800 *Res* 41: 182–190, 2007.
- 801 8. **Bartlett J, Hawley J, Morton J.** Carbohydrate availability and exercise training
802 adaptation: Too much of a good thing? *Eur J Sports Sci* 1–10, 2014.
- 803 9. **Bartlett J, Louhelainen J, Iqbal Z, Cochran A, Gibala M, Gregson W, Close G, Drust B,**
804 **Morton J.** Reduced carbohydrate availability enhances exercise-induced p53 signaling in
805 human skeletal muscle: implications for mitochondrial biogenesis. *Am J Physiol Regul Integr*
806 *Comp Physiol* 304: R450–R458, 2013.
- 807 10. **Bejma J, Ji LL.** Aging and acute exercise enhance free radical generation in rat skeletal
808 muscle. *J Appl Physiol* 87: 465–470, 1999.
- 809 11. **Bergström J, Hermansen L, Hultman E, Saltin B.** Diet, muscle glycogen and physical
810 performance. *Acta Physiol Scand* 71: 140–150, 1967.
- 811 12. **Bleakley C, Glasgow P, Phillips N, Hanna L, Callaghan M, Davison G, Hopkins T,**
812 **Delahunt E.** Management of acute soft tissue injury using Protection Rest Ice Compression
813 and Elevation Association of Chartered Physiotherapists in Sports and Exercise Medicine,
814 2010.
- 815 13. **Blomstrand E, Saltin B.** Effect of muscle glycogen on glucose, lactate and amino acid
816 metabolism during exercise and recovery in human subjects. *J Physiol* 514: 293–302, 1999.
- 817 14. **Bobeuf F, Labonte M, Dionne IJ, Khalil A.** Combined effect of antioxidant
818 supplementation and resistance training on oxidative stress markers, muscle and body
819 composition in an elderly population. *J Nutr Health Aging* 15: 883–889, 2011.
- 820 15. **Borzelleca JF.** Paracelsus: herald of modern toxicology. *Toxicol Sci* 53: 2–4, 2000.
- 821 16. **Broome CS, Kayani AC, Palomero J, Dillmann WH, Mestril R, Jackson MJ, McArdle A.**
822 Effect of lifelong overexpression of HSP70 in skeletal muscle on age-related oxidative stress
823 and adaptation after nondamaging contractile activity. *FASEB J* 20: 1549–1551, 2006.

17. **Brunmair B, Staniek K, Dorig J, Szocs Z, Stadlbauer K, Marian V, Gras F, Anderwald C, Nohl H, Waldhausl W, Furnsinn C.** Activation of PPAR- δ in isolated rat skeletal muscle switches fuel preference from glucose to fatty acids. *Diabetologia* 49: 2713–2722, 2006.
18. **Bruusgaard JC, Johansen IB, Egner IM, Rana ZA, Gundersen K.** Myonuclei acquired by overload exercise precede hypertrophy and are not lost on detraining. *Proc Natl Acad Sci U S A* 107: 15111–15116, 2010.
19. **Burd N, Dickinson J, Lemoine J, Carroll C, Sullivan B, Haus J, Jemiole B, Trappe S, Hughes G, Sanders C, Jr, Trappe T.** Effect of a cyclooxygenase-2 inhibitor on postexercise muscle protein synthesis in humans. *Am J Physiol Endocrinol Metab* 298: E354–E361, 2010.
20. **Burian A, Frangione V, Rovati S, Mautone G, Leuratti C, Vaccani A, Crevenna R, Keilani M, Burian B, Brunner M, Zeitlinger M.** An exploratory microdialysis study investigating the effect of repeated application of a diclofenac epolamine medicated plaster on prostaglandin concentrations in skeletal muscle after standardized physical exercise. *Br J Clin Pharmacol* 76: 880–887, 2013.
21. **Burke L, Hawley J, Wong S, Jeukendrup A.** Carbohydrates for training and competition. *J Sports Sci* 29: S17–S27, 2011.
22. **Calabrese EJ, Bachmann KA, Bailer AJ, Bolger PM, Borak J, Cai L, Cedergreen N, Cherian MG, Chiueh CC, Clarkson TW, Cook RR, Diamond DM, Doolittle DJ, Dorato MA, Duke SO, Feinendegen L, Gardner DE, Hart RW, Hastings KL, Hayes AW, Hoffmann GR, Ives JA, Jaworowski Z, Johnson TE, Jonas WB, Kaminski NE, Keller JG, Klaunig JE, Knudsen TB, Kozumbo WJ, Lettieri T, Liu SZ, Maisseu A, Maynard KI, Masoro EJ, McClellan RO, Mehendale HM, Mothersill C, Newlin DB, Nigg HN, Oehme FW, Phalen RF, Philbert MA, Rattan SI, Riviere JE, Rodricks J, Sapolsky RM, Scott BR, Seymour C, Sinclair DA, Smith-Sonneborn J, Snow ET, Spear L, Stevenson DE, Thomas Y, Tubiana M, Williams GM, Mattson MP.** Biological stress response terminology: Integrating the concepts of adaptive response and preconditioning stress within a hormetic dose-response framework. *Toxicol Appl Pharmacol* 222: 122–128, 2007.
23. **Camera D, West D, Burd N, Phillips S, Garnham A, Hawley J, Coffey V.** Low muscle glycogen concentration does not suppress the anabolic response to resistance exercise. *J Appl Physiol* 113: 206–214, 2012.
24. **Carroll C, O'Connor D, Steinmeyer R, Del Mundo J, McMullan D, Whitt J, Ramos J, Gonzales R.** The influence of acute resistance exercise on cyclooxygenase-1 and -2 activity and protein levels in human skeletal muscle. *Am J Physiol Regul Integr Comp Physiol* 305: R24–R30, 2013.
25. **Carvalho N, Puntel G, Correa P, Gubert P, Amaral G, Morais J, Royes L, da Rocha J, Soares F.** Protective effects of therapeutic cold and heat against the oxidative damage induced by a muscle strain injury in rats. *J Sports Sci* 28: 923–935, 2010.
26. **Chen HL, Nosaka K, Chen TC.** Muscle damage protection by low-intensity eccentric contractions remains for 2 weeks but not 3 weeks. *Eur J Appl Physiol* 112: 555–565, 2012.
27. **Chen HL, Nosaka K, Pearce AJ, Chen TC.** Two maximal isometric contractions attenuate the magnitude of eccentric exercise-induced muscle damage. *Appl Physiol Nutr Metab* 37: 680–689, 2012.
28. **Chen TC, Chen HL, Lin MJ, Chen CH, Pearce AJ, Nosaka K.** Effect of two maximal isometric contractions on eccentric exercise-induced muscle damage of the elbow flexors. *Eur J Appl Physiol* 113: 1545–1554, 2013.
29. **Chen TC, Chen HL, Lin MJ, Wu CJ, Nosaka K.** Potent protective effect conferred by four bouts of low-intensity eccentric exercise. *Med Sci Sports Exerc* 42: 1004–1012, 2010.

- 871 30. **Chen TC, Chen HL, Pearce AJ, Nosaka K.** Attenuation of eccentric exercise-induced
872 muscle damage by preconditioning exercises. *Med Sci Sports Exerc* 44: 2090–2098, 2012.
- 873 31. **Cheung K, Hume P, Maxwell L.** Delayed onset muscle soreness: treatment strategies and
874 performance factors. *Sports Med* 33: 145–164, 2003.
- 875 32. **Coyle E.** Physical activity as a metabolic stressor. *Am J Clin Nutr* 72: 512s–520s, 2000.
- 876 33. **Cumming KT, Paulsen G, Wernbom M, Ugelstad I, Raastad T.** Acute response and
877 subcellular movement of HSP27, α B-crystallin and HSP70 in human skeletal muscle after
878 blood-flow-restricted low-load resistance exercise. *Acta Physiol (Oxf)* 211: 634–646, 2014.
- 879 34. **Davies KJ, Quintanilha AT, Brooks GA, Packer L.** Free radicals and tissue damage
880 produced by exercise. *Biochem Biophys Res Commun* 107: 1198–1205, 1982.
- 881 35. **Davies P, Bailey PJ, Goldenberg MM, Ford-Hutchinson AW.** The role of arachidonic acid
882 oxygenation products in pain and inflammation. *Annual Rev Immunol* 2: 335–357, 1984.
- 883 36. **DeWitt DL, Meade EA, Smith WL.** PGH synthase isoenzyme selectivity: the potential for
884 safer nonsteroidal antiinflammatory drugs. *Am J Clin Nutr* 95: 40S–44s, 1993.
- 885 37. **Dodd SL, Gagnon BJ, Senf SM, Hain BA, Judge AR.** ROS-mediated activation of NF- κ B
886 and FOXO during muscle disuse. *Muscle Nerve* 41: 110–113, 2010.
- 887 38. **Dodd SL, Hain B, Senf SM, Judge AR.** Hsp27 inhibits IKK β -induced NF- κ B activity and
888 skeletal muscle atrophy. *FASEB J* 23: 3415–3423, 2009.
- 889 39. **Dousset E, Avela J, Ishikawa M, Kallio J, Kuitunen S, Kyröläinen H, Linnamo V, Komi PV.**
890 Bimodal recovery pattern in human skeletal muscle induced by exhaustive stretch-
891 shortening cycle exercise. *Med Sci Sports Exerc* 39: 453–460, 2007.
- 892 40. **Egan B, Zierath JR.** Exercise metabolism and the molecular regulation of skeletal muscle
893 adaptation. *Cell Metab* 17: 162–184, 2013.
- 894 41. **Enos R, Davis J, McClellan J, Murphy E.** Indomethacin in combination with exercise leads
895 to muscle and brain inflammation in mice. *J Interferon Cytokine Res* 33: 446–451, 2013.
- 896 42. **Flower RJ.** Studies on the mechanism of action of anti-inflammatory drugs. A paper in
897 honour of John Vane. *Thrombosis Research* 110: 259–263, 2003.
- 898 43. **Folkesson M, Mackey AL, Holm L, Kjaer M, Paulsen G, Raastad T, Henriksson J, Kadi F.**
899 Immunohistochemical changes in the expression of HSP27 in exercised human vastus
900 lateralis muscle. *Acta Physiol (Oxf)* 194: 215–222, 2008.
- 901 44. **Frohlich M, Faude O, Klein M, Pieter A, Emrich E, Meyer T.** Strength training adaptations
902 after cold water immersion. *J Strength Cond Res* 14: 2628–2633, 2014.
- 903 45. **Fry CS, Glynn EL, Drummond MJ, Timmerman KL, Fujita S, Abe T, Dhanani S, Volpi E,**
904 **Rasmussen BB.** Blood flow restriction exercise stimulates mTORC1 signaling and muscle
905 protein synthesis in older men. *J Appl Physiol* 108: 1199–1209, 2010.
- 906 46. **Fu FH, Cen HW, Eston RG.** The effects of cryotherapy on muscle damage in rats
907 subjected to endurance training. *Scand J Med Sci Sports* 7: 358–362, 1997.
- 908 47. **Fujita S, Abe T, Drummond MJ, Cadenas JG, Dreyer HC, Sato Y, Volpi E, Rasmussen BB.**
909 Blood flow restriction during low-intensity resistance exercise increases S6K1
910 phosphorylation and muscle protein synthesis. *J Appl Physiol* 103: 903–910, 2007.
- 911 48. **Garramone RR, Jr., Winters RM, Das DK, Deckers PJ.** Reduction of skeletal muscle injury
912 through stress conditioning using the heat-shock response. *Plast Reconstr Surg* 93: 1242–
913 1247, 1994.
- 914 49. **Gehrig SM, van der Poel C, Sayer TA, Schertzer JD, Henstridge DC, Church JE, Lamon S,**
915 **Russell AP, Davies KE, Febbraio MA, Lynch GS.** Hsp72 preserves muscle function and slows
916 progression of severe muscular dystrophy. *Nature* 484: 394–398, 2012.

50. Gejl K, Hvid L, Frandsen U, Jensen K, Sahlin K, Ørtenblad N. Muscle glycogen content modifies SR Ca^{2+} release rate in elite endurance athletes. *Med Sci Sports Exerc* 46: 496–505, 2014.
51. Gerhart-Hines Z, Rodgers JT, Bare O, Lerin C, Kim SH, Mostoslavsky R, Alt FW, Wu Z, Puigserver P. Metabolic control of muscle mitochondrial function and fatty acid oxidation through SIRT1/PGC-1 α . *EMBO J* 26: 1913–1923, 2007.
52. Giombini A, Giovannini V, Di Cesare A, Pacetti P, Ichinoseki-Sekine N, Shiraishi M, Naito H, Maffulli N. Hyperthermia induced by microwave diathermy in the management of muscle and tendon injuries. *Br Med Bull* 83: 379–396, 2007.
53. Gomez-Cabrera MC, Borrás C, Pallardo FV, Sastre J, Ji LL, Vina J. Decreasing xanthine oxidase-mediated oxidative stress prevents useful cellular adaptations to exercise in rats. *J Physiol* 567: 113–120, 2005.
54. Gomez-Cabrera MC, Domenech E, Romagnoli M, Arduini A, Borrás C, Pallardo FV, Sastre J, Vina J. Oral administration of vitamin C decreases muscle mitochondrial biogenesis and hampers training-induced adaptations in endurance performance. *Am J Clin Nutr* 87: 142–149, 2008.
55. Goto K, Okuyama R, Sugiyama H, Honda M, Kobayashi T, Uehara K, Akema T, Sugiura T, Yamada S, Ohira Y, Yoshioka T. Effects of heat stress and mechanical stretch on protein expression in cultured skeletal muscle cells. *Pflugers Arch* 447: 247–253, 2003.
56. Granado M, Martin AI, Villanua MA, Lopez-Calderon A. Experimental arthritis inhibits the insulin-like growth factor-I axis and induces muscle wasting through cyclooxygenase-2 activation. *Am J Physiol Endocrinol Metab* 292: E1656–E1665, 2007.
57. Griswold DE, Adams JL. Constitutive cyclooxygenase (COX-1) and inducible cyclooxygenase (COX-2): rationale for selective inhibition and progress to date. *Medicinal Res Rev* 16: 181–206, 1996.
58. Grounds MD. Phagocytosis of necrotic muscle in muscle isografts is influenced by the strain, age, and sex of host mice. *J Pathol* 153: 71–82, 1987.
59. Gump B, McMullan D, Cauthon D, Whitt J, Del Mundo J, Letham T, Kim P, Friedlander G, Pingel J, Langberg H, Carroll C. Short-term acetaminophen consumption enhances the exercise-induced increase in Achilles peritendinous IL-6 in humans. *J Appl Physiol* 115: 929–936, 2013.
60. Gundermann DM, Walker DK, Reidy PT, Borack MS, Dickinson JM, Volpi E, Rasmussen BB. Activation of mTORC1 signaling and protein synthesis in human muscle following blood flow restriction exercise is inhibited by rapamycin. *Am J Physiol Endocrinol Metab* 306: E1198–E1204, 2014.
61. Gwag T, Park K, Kim E, Son C, Park J, Nikawa T, Choi I. Inhibition of C2C12 myotube atrophy by a novel HSP70 inducer, celastrol, via activation of Akt1 and ERK1/2 pathways. *Arch Biochem Biophys* 537: 21–30, 2013.
62. Haddad F, Zaldivar F, Cooper DM, Adams GR. IL-6-induced skeletal muscle atrophy. *J Appl Physiol* 98: 911–917, 2005.
63. Haddix TL, Pohlman TH, Noel RF, Sato TT, Boyle EM, Jr., Verrier ED. Hypothermia inhibits human E-selectin transcription. *J Surg Res* 64: 176–183, 1996.
64. Hammers DW, Rybalko V, Merscham-Banda M, Hsieh PL, Suggs LJ, Farrar RP. Anti-inflammatory macrophages improve skeletal muscle recovery from ischemia/reperfusion. *J Appl Physiol* in press, 2014. doi: 10.1152/jappphysiol.00313.2014

- 962 65. **Hansen A, Fischer C, Plomgaard P, Andersen J, Saltin B, Pedersen B.** Skeletal muscle
 963 adaptation: training twice every second day vs. training once daily. *J Appl Physiol* 98: 93–99,
 964 2005.
- 965 66. **Hardie D.** AMP-activated protein kinase: maintaining energy homeostasis at the cellular
 966 and whole-body levels. *Annual Rev Nutr* 34: 31–55, 2014.
- 967 67. **Hardie D, Ross F, Hawley S.** AMPK: a nutrient and energy sensor that maintains energy
 968 homeostasis. *Nat Rev Mol Cell Biol* 13: 251–262, 2012.
- 969 68. **Hatade T, Takeuchi K, Fujita N, Arakawa T, Miki A.** Effect of heat stress soon after
 970 muscle injury on the expression of MyoD and myogenin during regeneration process. *J*
 971 *Musculoskelet Neuronal Interact* 14: 325–333, 2014.
- 972 69. **Hawley J, Morton J.** Ramping up the signal: promoting endurance training adaptation in
 973 skeletal muscle by nutritional manipulation. *Clin Exp Pharmacol Physiol* 41: 608–613, 2014.
- 974 70. **Higashida K, Kim SH, Higuchi M, Holloszy JO, Han DH.** Normal adaptations to exercise
 975 despite protection against oxidative stress. *Am J Physiol Endocrinol Metab* 301: E779–E784,
 976 2011.
- 977 71. **Hirunsai M, Srikuea R, Yimlamai T.** Heat stress promotes extracellular matrix
 978 remodelling via TGF- β 1 and MMP-2/TIMP-2 modulation in tenotomised soleus and plantaris
 979 muscles. *Int J Hyperthermia* 1–13, 2015.
- 980 72. **Howarth K, Phillips S, MacDonald M, Richards D, Moreau N, Gibala M.** Effect of
 981 glycogen availability on human skeletal muscle protein turnover during exercise and
 982 recovery. *J Appl Physiol* 109: 431–438, 2010.
- 983 73. **Howatson G, van Someren KA.** Evidence of a contralateral repeated bout effect after
 984 maximal eccentric contractions. *Eur J Appl Physiol* 101: 207–214, 2007.
- 985 74. **Hubbard TJ, Aronson SL, Denegar CR.** Does cryotherapy hasten return to participation?
 986 A systematic review. *J Athl Train* 39: 88–94, 2004.
- 987 75. **Hulston C, Venables M, Mann C, Martin C, Philp A, Baar K, Jeukendrup A.** Training with
 988 low muscle glycogen enhances fat metabolism in well-trained cyclists. *Med Sci Sports Exerc*
 989 42: 2046–2055, 2010.
- 990 76. **Hurme T, Rantanen J, Kaliomo H.** Effects of early cryotherapy in experimental skeletal
 991 muscle injury. *Scand J Med Sci Sports* 3: 46–51, 1993.
- 992 77. **Iguchi M, Shields RK.** Prior heat stress effects fatigue recovery of the elbow flexor
 993 muscles. *Muscle Nerve* 44: 115–125, 2011.
- 994 78. **Inamasu J, Suga S, Sato S, Horiguchi T, Akaji K, Mayanagi K, Kawase T.** Intra-ischemic
 995 hypothermia attenuates intercellular adhesion molecule-1 (ICAM-1) and migration of
 996 neutrophil. *Neurol Res* 23: 105–111, 2001.
- 997 79. **Irrcher I, Ljubicic V, Hood D.** Interactions between ROS and AMP kinase activity in the
 998 regulation of PGC-1 α transcription in skeletal muscle cells. *Am J Physiol Cell Physiol* 296:
 999 C116–C123, 2009.
- 1000 80. **Irrcher I, Ljubicic V, Kirwan AF, Hood DA.** AMP-activated protein kinase-regulated
 1001 activation of the PGC-1 α promoter in skeletal muscle cells. *PLoS One* 3: e3614, 2008.
- 1002 81. **Ito N, Ruegg UT, Kudo A, Miyagoe-Suzuki Y, Takeda S.** Activation of calcium signaling
 1003 through Trpv1 by nNOS and peroxynitrite as a key trigger of skeletal muscle hypertrophy.
 1004 *Nature Med* 19: 101–106, 2013.
- 1005 82. **Iversen E, Rostad V.** Low-load ischemic exercise-induced rhabdomyolysis. *Clin J Sport*
 1006 *Med* 20: 218–219, 2010.

- 1007 83. Jager S, Handschin C, St-Pierre J, Spiegelman BM. AMP-activated protein kinase (AMPK)
1008 action in skeletal muscle via direct phosphorylation of PGC-1 α . *Proc Natl Acad Sci U S A* 104:
1009 12017–12022, 2007.
- 1010 84. Jang YC, Liu Y, Hayworth CR, Bhattacharya A, Lustgarten MS, Muller FL, Chaudhuri A,
1011 Qi W, Li Y, Huang JY, Verdin E, Richardson A, Van Remmen H. Dietary restriction attenuates
1012 age-associated muscle atrophy by lowering oxidative stress in mice even in complete
1013 absence of CuZnSOD. *Aging Cell* 11: 770–782, 2012.
- 1014 85. Jankowski C, Gozansky W, MacLean P, Shulman B, Wolfe P, Schwartz R, Kohrt W. N-
1015 acetyl-4-aminophenol and musculoskeletal adaptations to resistance exercise training. *Eur J*
1016 *Appl Physiol* 113: 1127–1136, 2013.
- 1017 86. Jayaraman RC, Reid RW, Foley JM, Prior BM, Dudley GA, Weingand KW, Meyer RA. MRI
1018 evaluation of topical heat and static stretching as therapeutic modalities for the treatment
1019 of eccentric exercise-induced muscle damage. *Eur J Appl Physiol* 93: 30–38, 2004.
- 1020 87. Jiang B, Xiao W, Shi Y, Liu M, Xiao X. Heat shock pretreatment inhibited the release of
1021 Smac/DIABLO from mitochondria and apoptosis induced by hydrogen peroxide in
1022 cardiomyocytes and C2C12 myogenic cells. *Cell Stress Chaperones* 10: 252–262, 2005.
- 1023 88. Jung MH, Song MC, Bae K, Kim HS, Kim SH, Sung SH, Ye SK, Lee KH, Yun YP, Kim TJ.
1024 Sauchinone attenuates oxidative stress-induced skeletal muscle myoblast damage through
1025 the down-regulation of ceramide. *Biol Pharm Bull* 34: 575–579, 2011.
- 1026 89. Kabakov AE, Budagova KR, Latchman DS, Kampinga HH. Stressful preconditioning and
1027 HSP70 overexpression attenuate proteotoxicity of cellular ATP depletion. *Am J Physiol Cell*
1028 *Physiol* 283: C521–C534, 2002.
- 1029 90. Kakigi R, Naito H, Ogura Y, Kobayashi H, Saga N, Ichinoseki-Sekine N, Yoshihara T,
1030 Katamoto S. Heat stress enhances mTOR signaling after resistance exercise in human
1031 skeletal muscle. *J Physiol Sci* 61: 131–140, 2011.
- 1032 91. Kang C, O'Moore KM, Dickman JR, Ji LL. Exercise activation of muscle peroxisome
1033 proliferator-activated receptor- γ coactivator-1 α signaling is redox sensitive. *Free Radic Biol*
1034 *Med* 47: 1394–1400, 2009.
- 1035 92. Karamouzis I, Christoulas K, Grekas D, Giannoulis K, Vamvakoudis E, Mandroukas K.
1036 The response of muscle interstitial F₂-isoprostane (8-iso-PGF_{2 α}) during dynamic muscle
1037 contractions in humans. *Prostaglandins Leukot Essent Fatty Acid* 71: 87–90, 2004.
- 1038 93. Karamouzis M, Karamouzis I, Vamvakoudis E, Ampatzidis G, Christoulas K,
1039 Angelopoulou N, Mandroukas K. The response of muscle interstitial prostaglandin E₂ (PGE₂),
1040 prostacyclin I₂ (PGI₂) and thromboxane A₂(TXA₂) levels during incremental dynamic exercise
1041 in humans determined by in vivo microdialysis. *Prostaglandins Leukot Essent Fatty Acid* 64:
1042 259–263, 2001.
- 1043 94. Karlsson J, Saltin B. Diet, muscle glycogen, and endurance performance. *J Appl Physiol*
1044 31: 203–206, 1971.
- 1045 95. Khoshkharesh F, Siahkuhain M, Fisher G, Nakhostin-Rooh B. Influence of a low-dose
1046 cox-2 inhibitor drug on exercise-induced inflammation, muscle damage and lipid
1047 peroxidation. *Biol Sport* 30: 61–65, 2013.
- 1048 96. Kojima A, Goto K, Morioka S, Naito T, Akema T, Fujiya H, Sugiura T, Ohira Y, Beppu M,
1049 Aoki H, Yoshioka T. Heat stress facilitates the regeneration of injured skeletal muscle in rats.
1050 *J Orthop Sci* 12: 74–82, 2007.
- 1051 97. Korotkova M, Lundberg I. The skeletal muscle arachidonic acid cascade in health and
1052 inflammatory disease. *Nature Rev Rheum* 10: 295–303, 2014.

- 1053 98. **Koya T, Nishizawa S, Ohno Y, Goto A, Ikuta A, Suzuki M, Ohira T, Egawa T, Nakai A,**
1054 **Sugiura T, Ohira Y, Yoshioka T, Beppu M, Goto K.** Heat shock transcription factor 1-
1055 deficiency attenuates overloading-associated hypertrophy of mouse soleus muscle. *PLoS*
1056 *One* 8: e77788, 2013.
- 1057 99. **Krentz J, Quest B, Farthing J, Quest D, Chilibeck P.** The effects of ibuprofen on muscle
1058 hypertrophy, strength, and soreness during resistance training. *Appl Physiol Nutr Metab* 33:
1059 470–475, 2008.
- 1060 100. **Lapointe BM, Fremont P, Cote CH.** Adaptation to lengthening contractions is
1061 independent of voluntary muscle recruitment but relies on inflammation. *Am J Physiol Regul*
1062 *Integr Comp Physiol* 282: R323–R329, 2002.
- 1063 101. **Lavender AP, Nosaka K.** A light load eccentric exercise confers protection against a
1064 subsequent bout of more demanding eccentric exercise. *J Sci Med Sport* 11: 291–298, 2008.
- 1065 102. **Lee H, Natsui H, Akimoto T, Yanagi K, Ohshima N, Kono I.** Effects of cryotherapy
1066 after contusion using real-time intravital microscopy. *Med Sci Sports Exerc* 37: 1093–1098,
1067 2005.
- 1068 103. **Lee IM, Hsieh CC, Paffenbarger RS, Jr.** Exercise intensity and longevity in men. The
1069 Harvard Alumni Health Study. *JAMA* 273: 1179–1184, 1995.
- 1070 104. **Lehmann JF, Dundore DE, Esselman PC, Nelp WB.** Microwave diathermy: effects on
1071 experimental muscle hematoma resolution. *Arch Phys Med Rehabil* 64: 127–129, 1983.
- 1072 105. **Liu CC, Lin CH, Lin CY, Lee CC, Lin MT, Wen HC.** Transgenic overexpression of heat
1073 shock protein 72 in mouse muscle protects against exhaustive exercise-induced skeletal
1074 muscle damage. *J Formos Med Assoc* 112: 24–30, 2013.
- 1075 106. **Loell I, Alemo Munters L, Pandya J, Zong M, Alexanderson H, Fasth AE, Stahl**
1076 **Hallengren C, Radmark O, Lundberg IE, Jakobsson PJ, Korotkova M.** Activated LTB4
1077 pathway in muscle tissue of patients with polymyositis or dermatomyositis. *Ann Rheum Dis*
1078 72: 293–299, 2013.
- 1079 107. **Loenneke JP, Abe T.** Does blood flow restricted exercise result in prolonged torque
1080 decrements and muscle damage? *Eur J Appl Physiol* 112: 3445–3446, 2012.
- 1081 108. **Loenneke JP, Abe T, Wilson JM, Thiebaud RS, Fahs CA, Rossow LM, Bembien MG.**
1082 Blood flow restriction: an evidence based progressive model (Review). *Acta Physiol Hung* 99:
1083 235–250, 2012.
- 1084 109. **Loenneke JP, Fahs CA, Rossow LM, Thiebaud RS, Mattocks KT, Abe T, Bembien MG.**
1085 Blood flow restriction pressure recommendations: a tale of two cuffs. *Front Physiol* 4: 249,
1086 2013.
- 1087 110. **Loenneke JP, Fahs CA, Thiebaud RS, Rossow LM, Abe T, Ye X, Kim D, Bembien MG.**
1088 The acute muscle swelling effects of blood flow restriction. *Acta Physiol Hung* 99: 400–410,
1089 2012.
- 1090 111. **Loenneke JP, Kearney ML, Thrower AD, Collins S, Pujol TJ.** The acute response of
1091 practical occlusion in the knee extensors. *J Strength Cond Res* 24: 2831–2834, 2010.
- 1092 112. **Loenneke JP, Thiebaud RS, Fahs CA, Rossow LM, Abe T, Bembien MG.** Blood flow
1093 restriction does not result in prolonged decrements in torque. *Eur J Appl Physiol* 113: 923–
1094 931, 2013.
- 1095 113. **Loenneke JP, Wilson GJ, Wilson JM.** A mechanistic approach to blood flow occlusion.
1096 *Int J Sports Med* 31: 1–4, 2010.
- 1097 114. **Loenneke JP, Wilson JM, Marin PJ, Zourdos MC, Bembien MG.** Low intensity blood
1098 flow restriction training: a meta-analysis. *Eur J Appl Physiol* 112: 1849–1859, 2012.

- 1099 115. **Lu H, Huang D, Ransohoff RM, Zhou L.** Acute skeletal muscle injury: CCL2 expression
1100 by both monocytes and injured muscle is required for repair. *FASEB J* 2011.
- 1101 116. **Lu H, Huang D, Saederup N, Charo IF, Ransohoff RM, Zhou L.** Macrophages recruited
1102 via CCR2 produce insulin-like growth factor-1 to repair acute skeletal muscle injury. *FASEB J*
1103 25: 358–369, 2011.
- 1104 117. **Mackey A, Kjaer M, Dandanell S, Mikkelsen K, Holm L, Døssing S, Fawzi K, Koskinen**
1105 **S, Jensen C, der H, Langberg H.** The influence of anti-inflammatory medication on exercise-
1106 induced myogenic precursor cell responses in humans. *J Appl Physiol* 103: 425–431, 2007.
- 1107 118. **Maglara AA, Vasilaki A, Jackson MJ, McArdle A.** Damage to developing mouse
1108 skeletal muscle myotubes in culture: protective effect of heat shock proteins. *J Physiol* 548:
1109 837–846, 2003.
- 1110 119. **Makanae Y, Kawada S, Sasaki K, Nakazato K, Ishii N.** Vitamin C administration
1111 attenuates overload-induced skeletal muscle hypertrophy in rats. *Acta Physiol* 208: 57–65,
1112 2013.
- 1113 120. **Mankowski RT, Anton SD, Buford TW, Leeuwenburgh C.** Dietary antioxidants as
1114 modifiers of physiologic adaptations to exercise. *Med Sci Sports Exerc* in press: 2015.
- 1115 121. **Marber MS, Latchman DS, Walker JM, Yellon DM.** Cardiac stress protein elevation
1116 24 hours after brief ischemia or heat stress is associated with resistance to myocardial
1117 infarction. *Circulation* 88: 1264–1272, 1993.
- 1118 122. **Markworth J, Vella L, Figueiredo V, Cameron-Smith D.** Ibuprofen treatment blunts
1119 early translational signaling responses in human skeletal muscle following resistance
1120 exercise. *J Appl Physiol* 117: 20–28, 2014.
- 1121 123. **Markworth J, Vella L, Lingard B, Tull D, Rupasinghe T, Sinclair A, Maddipati K,**
1122 **Cameron-Smith D.** Human inflammatory and resolving lipid mediator responses to
1123 resistance exercise and ibuprofen treatment. *Am J Physiol Regul Integr Comp Physiol* 305:
1124 R1281–R1296, 2013.
- 1125 124. **Markworth JF, Cameron-Smith D.** Arachidonic acid supplementation enhances in
1126 vitro skeletal muscle cell growth via a COX-2-dependent pathway. *Am J Physiol Cell Physiol*
1127 304: C56–C67, 2013.
- 1128 125. **Markworth JF, Cameron-Smith D.** Prostaglandin $F_{2\alpha}$ stimulates PI3K/ERK/mTOR
1129 signaling and skeletal myotube hypertrophy. *Am J Physiol Cell Physiol* 300: C671–C682,
1130 2011.
- 1131 126. **Marshall RJ, Scott KC, Hill RC, Lewis DD, Sundstrom D, Jones GL, Harper J.**
1132 Supplemental vitamin C appears to slow racing greyhounds. *J Nutr* 132: 1616S–1621S, 2002.
- 1133 127. **Mattson MP.** Hormesis defined. *Ageing Res Rev* 7: 1–7, 2008.
- 1134 128. **McBride A, Ghilagaber S, Nikolaev A, Hardie D.** The glycogen-binding domain on the
1135 AMPK β subunit allows the kinase to act as a glycogen sensor. *Cell Metab* 9: 23–34, 2009.
- 1136 129. **McCarthy DO, Whitney P, Hitt A, Al-Majid S.** Indomethacin and ibuprofen preserve
1137 gastrocnemius muscle mass in mice bearing the colon-26 adenocarcinoma. *Res Nurs Health*
1138 27: 174–184, 2004.
- 1139 130. **McConell GK, Ng GPY, Phillips M, Ruan Z, Macaulay SL, Wadley GD.** Central role of
1140 nitric oxide synthase in AICAR and caffeine induced mitochondrial biogenesis in L6
1141 myocytes. *J Appl Physiol* 108: 589–595, 2010.
- 1142 131. **McHugh MP, Connolly DA, Eston RG, Gleim GW.** Exercise-induced muscle damage
1143 and potential mechanisms for the repeated bout effect. *Sports Med* 27: 157–170, 1999.
- 1144 132. **Meeusen R, Lievens P.** The use of cryotherapy in sports injuries. *Sports Med* 3: 398–
1145 414, 1986.

- 1146 133. **Merrick MA, McBrier NM.** Progression of secondary injury after musculoskeletal
1147 trauma—a window of opportunity? *J Sport Rehabil* 19: 380–388, 2010.
- 1148 134. **Merrick MA, Rankin JM, Andres FA, Hinman CL.** A preliminary examination of
1149 cryotherapy and secondary injury in skeletal muscle. *Med Sci Sports Exerc* 31: 1516–1521,
1150 1999.
- 1151 135. **Mikkelsen U, Helmark I, Kær M, Langberg H.** Prostaglandin synthesis can be
1152 inhibited locally by infusion of NSAIDS through microdialysis catheters in human skeletal
1153 muscle. *J Appl Physiol* 104: 534–537, 2008.
- 1154 136. **Mikkelsen U, Paulsen G, Schjerling P, Helmark I, Langberg H, Kjær M, Heinemeier K.**
1155 The heat shock protein response following eccentric exercise in human skeletal muscle is
1156 unaffected by local NSAID infusion. *Eur J Appl Physiol* 113: 1883–1893, 2013.
- 1157 137. **Mikkelsen UR, Langberg H, Helmark IC, Skovgaard D, Andersen LL, Kjær M, Mackey**
1158 **AL.** Local NSAID infusion inhibits satellite cell proliferation in human skeletal muscle after
1159 eccentric exercise. *J Appl Physiol* 107: 1600–1611, 2009.
- 1160 138. **Mikkelsen UR, Schjerling P, Helmark IC, Reitelseder S, Holm L, Skovgaard D,**
1161 **Langberg H, Kjær M, Heinemeier KM.** Local NSAID infusion does not affect protein synthesis
1162 and gene expression in human muscle after eccentric exercise. *Scand J Med Sci Sports* 21:
1163 630–644, 2011.
- 1164 139. **Morita I.** Distinct functions of COX-1 and COX-2. *Prostaglandins Other Lipid Mediat*
1165 68–69: 165–175, 2002.
- 1166 140. **Morton J, Croft L, Bartlett J, MacLaren D, Reilly T, Evans L, McArdle A, Drust B.**
1167 Reduced carbohydrate availability does not modulate training-induced heat shock protein
1168 adaptations but does upregulate oxidative enzyme activity in human skeletal muscle. *J Appl*
1169 *Physiol* 106: 1513–1521, 2009.
- 1170 141. **Murry CE, Jennings RB, Reimer KA.** Preconditioning with ischemia: a delay of lethal
1171 cell injury in ischemic myocardium. *Circulation* 74: 1124–1136, 1986.
- 1172 142. **Naito H, Powers SK, Demirel HA, Sugiura T, Dodd SL, Aoki J.** Heat stress attenuates
1173 skeletal muscle atrophy in hindlimb-unweighted rats. *J Appl Physiol* 88: 359–363, 2000.
- 1174 143. **Nguyen HX, Tidball JG.** Null mutation of gp91phox reduces muscle membrane lysis
1175 during muscle inflammation in mice. *J Physiol* 553: 833–841, 2003.
- 1176 144. **Nielsen J, Cheng AJ, Ørtenblad N, Westerblad H.** Subcellular distribution of glycogen
1177 and decreased tetanic Ca²⁺ in fatigued single intact mouse muscle fibres. *J Physiol* 592:
1178 2003–2012, 2014.
- 1179 145. **Nielsen JL, Aagaard P, Bech RD, Nygaard T, Hvid LG, Wernbom M, Suetta C,**
1180 **Frandsen U.** Proliferation of myogenic stem cells in human skeletal muscle in response to
1181 low-load resistance training with blood flow restriction. *J Physiol* 590: 4351–4361, 2012.
- 1182 146. **Nieman D, Henson D, Dumke C, Oley K, McAnulty S, Davis J, Murphy E, Utter A, Lind**
1183 **R, McAnulty L, Morrow J.** Ibuprofen use, endotoxemia, inflammation, and plasma cytokines
1184 during ultramarathon competition. *Brain Behav Immun* 20: 578–584, 2006.
- 1185 147. **Nieman DC, Johanssen LM, Lee JW, Arabatzis K.** Infectious episodes in runners
1186 before and after the Los Angeles Marathon. *J Sports Med Phys Fitness* 30: 316–328, 1990.
- 1187 148. **Nishizawa S, Koya T, Ohno Y, Goto A, Ikuita A, Suzuki M, Ohira T, Egawa T, Nakai A,**
1188 **Sugiura T, Ohira Y, Yoshioka T, Beppu M, Goto K.** Regeneration of injured skeletal muscle in
1189 heat shock transcription factor 1-null mice. *Physiol Rep* 1: e00071, 2013.
- 1190 149. **Nosaka K, Muthalib M, Lavender A, Laursen PB.** Attenuation of muscle damage by
1191 preconditioning with muscle hyperthermia 1-day prior to eccentric exercise. *Eur J Appl*
1192 *Physiol* 99: 183–192, 2007.

- 1193 150. **Nosaka K, Sakamoto K, Newton M, Sacco P.** How long does the protective effect on
1194 eccentric exercise-induced muscle damage last? *Med Sci Sports Exerc* 33: 1490–1495, 2001.
- 1195 151. **Nosaka K, Sakamoto K, Newton M, Sacco P.** Influence of pre-exercise muscle
1196 temperature on responses to eccentric exercise. *J Athl Train* 39: 132–137, 2004.
- 1197 152. **Novak ML, Billich W, Smith SM, Sukhija KB, McLoughlin TJ, Hornberger TA, Koh TJ.**
1198 COX-2 inhibitor reduces skeletal muscle hypertrophy in mice. *Am J Physiol Regul Integr*
1199 *Comp Physiol* 296: R1132–R1139, 2009.
- 1200 153. **Ohnishi N, Yamane M, Uchiyama N, Shirasawa S, Kosaka M, Shiono H, Okada T.**
1201 Adaptive changes in muscular performance and circulation by resistance training with
1202 regular cold water immersion. *J Therm Biol* 29: 839–483, 2004.
- 1203 154. **Oishi Y, Hayashida M, Tsukiashi S, Taniguchi K, Kami K, Roy RR, Ohira Y.** Heat stress
1204 increases myonuclear number and fiber size via satellite cell activation in rat regenerating
1205 soleus fibers. *J Appl Physiol* 107: 1612–1621, 2009.
- 1206 155. **Ojuka EO, Jones TE, Han DH, Chen M, Holloszy JO.** Raising Ca^{2+} in L6 myotubes
1207 mimics effects of exercise on mitochondrial biogenesis in muscle. *FASEB J* 17: 675–681,
1208 2003.
- 1209 156. **Osterman AL, Heppenstall RB, Sapega AA, Katz M, Chance B, Sokolow D.** Muscle
1210 ischemia and hypothermia: a bioenergetic study using ^{31}P nuclear magnetic resonance
1211 spectroscopy. *J Trauma* 24: 811–817, 1984.
- 1212 157. **Paulsen G, Cumming KT, Holden G, Hallen J, Ronnestad BR, Sveen O, Skaug A, Paur**
1213 **I, Bastani NE, Ostgaard HN, Buer C, Midttun M, Freuchen F, Wiig H, Ulseth ET, Garthe I,**
1214 **Blomhoff R, Benestad HB, Raastad T.** Vitamin C and E supplementation hampers cellular
1215 adaptation to endurance training in humans: a double-blind, randomised, controlled trial. *J*
1216 *Physiol* 592: 1887–1901, 2014.
- 1217 158. **Paulsen G, Egner IM, Drange M, Langberg H, Benestad HB, Fjeld JG, Hallén J,**
1218 **Raastad T.** A COX-2 inhibitor reduces muscle soreness, but does not influence recovery and
1219 adaptation after eccentric exercise. *Scand J Med Sci Sports* 20: 1–13, 2010.
- 1220 159. **Paulsen G, Hamarsland H, Cumming KT, Johansen RE, Hulmi JJ, Borsheim E, Wiig H,**
1221 **Garthe I, Raastad T.** Vitamin C and E supplementation alters protein signalling after a
1222 strength training session, but not muscle growth during 10 weeks of training. *J Physiol* 592:
1223 5391–5408, 2014.
- 1224 160. **Pelosi L, Giacinti C, Nardis C, Borsellino G, Rizzuto E, Nicoletti C, Wannenes F,**
1225 **Battistini L, Rosenthal N, Molinaro M, Musaro A.** Local expression of IGF-1 accelerates
1226 muscle regeneration by rapidly modulating inflammatory cytokines and chemokines. *FASEB*
1227 *J* 21: 1393–1402, 2007.
- 1228 161. **Perry CG, Lally J, Holloway GP, Heigenhauser GJ, Bonen A, Spriet LL.** Repeated
1229 transient mRNA bursts precede increases in transcriptional and mitochondrial proteins
1230 during training in human skeletal muscle. *J Physiol* 588: 4795–4810, 2010.
- 1231 162. **Peternelj TT, Coombes JS.** Antioxidant supplementation during exercise training:
1232 beneficial or detrimental? *Sports Med* 41: 1043–1069, 2011.
- 1233 163. **Petersen AC, McKenna MJ, Medved I, Murphy KT, Brown MJ, Della Gatta P,**
1234 **Cameron-Smith D.** Infusion with the antioxidant N-acetylcysteine attenuates early adaptive
1235 responses to exercise in human skeletal muscle. *Acta physiologica* 204: 382–392, 2012.
- 1236 164. **Petersen S, Beyer N, Hansen M, Holm L, Aagaard P, Mackey A, Kjaer M.**
1237 Nonsteroidal anti-inflammatory drug or glucosamine reduced pain and improved muscle
1238 strength with resistance training in a randomized controlled trial of knee osteoarthritis
1239 patients. *Arch Phys Med Rehabil* 92: 1185–1193, 2011.

- 1240 165. **Petersen S, Miller B, Hansen M, Kjaer M, Holm L.** Exercise and NSAIDs: effect on
1241 muscle protein synthesis in patients with knee osteoarthritis. *Med Sci Sports Exerc* 43: 425–
1242 431, 2011.
- 1243 166. **Piestun Y, Harel M, Barak M, Yahav S, Halevy O.** Thermal manipulations in late-term
1244 chick embryos have immediate and longer term effects on myoblast proliferation and
1245 skeletal muscle hypertrophy. *J Appl Physiol* 106: 233–240, 2009.
- 1246 167. **Pizza FX, Cavender D, Stockard A, Baylies H, Beighle A.** Anti-inflammatory doses of
1247 ibuprofen: effect on neutrophils and exercise-induced muscle injury. *Int J Sports Med* 20:
1248 98–102, 1999.
- 1249 168. **Pizza FX, Peterson JM, Baas JH, Koh TJ.** Neutrophils contribute to muscle injury and
1250 impair its resolution after lengthening contractions in mice. *J Physiol* 562: 899–913, 2005.
- 1251 169. **Plaisance I, Morandi C, Murigande C, Brink M.** TNF- α increases protein content in
1252 C2C12 and primary myotubes by enhancing protein translation via the TNF-R1, PI3-kinase
1253 and MEK. *Am J Physiol Endocrinol Metab* 294: E241–E250, 2008.
- 1254 170. **Powers S, Smuder A, Judge A.** Oxidative stress and disuse muscle atrophy: cause or
1255 consequence? *Curr Opin Clin Nutr Metab Care* 15: 240–245, 2012.
- 1256 171. **Powers SK, Nelson WB, Hudson MB.** Exercise-induced oxidative stress in humans:
1257 cause and consequences. *Free Radic Biol Med* 51: 942–950, 2011.
- 1258 172. **Psilander N, Frank P, Flockhart M, Sahlin K.** Exercise with low glycogen increases
1259 PGC-1 α gene expression in human skeletal muscle. *Eur J Appl Physiol* 113: 951–963, 2013.
- 1260 173. **Puigserver P, Spiegelman BM.** Peroxisome proliferator-activated receptor- γ
1261 coactivator 1 α (PGC-1 α): transcriptional coactivator and metabolic regulator. *Endocr Rev*
1262 24: 78–90, 2003.
- 1263 174. **Puntel GO, Carvalho NR, Amaral GP, Lobato LD, Silveira SO, Daubermann MF,**
1264 **Barbosa NV, Rocha JB, Soares FA.** Therapeutic cold: An effective kind to modulate the
1265 oxidative damage resulting of a skeletal muscle contusion. *Free Radic Res* 45: 125–138,
1266 2011.
- 1267 175. **Radak Z, Chung HY, Goto S.** Exercise and hormesis: oxidative stress-related
1268 adaptation for successful aging. *Biogerontology* 6: 71–75, 2005.
- 1269 176. **Rieu I, Magne H, Savary-Auzeloux I, Averous J, Bos C, Peyron MA, Combaret L,**
1270 **Dardevet D.** Reduction of low grade inflammation restores blunting of postprandial muscle
1271 anabolism and limits sarcopenia in old rats. *J Physiol* 587: 5483–5492, 2009.
- 1272 177. **Ristow M, Zarse K, Oberbach A, Klötting N, Birringer M, Kiehntopf M, Stumvoll M,**
1273 **Kahn CR, Bluher M.** Antioxidants prevent health-promoting effects of physical exercise in
1274 humans. *Proc Natl Acad Sci U S A* 106: 8665–8670, 2009.
- 1275 178. **Roberts LA, Beattie K, Close GL, Morton JP.** Vitamin C consumption does not impair
1276 training-induced improvements in exercise performance. *Int J Sports Physiol Perf* 6: 58–69,
1277 2011.
- 1278 179. **Roberts LA, Raastad T, Cameron-Smith D, Coombes JS, Peake JM.** Cold water
1279 immersion reduces chronic resistance training-induced adaptation. *Med Sci Sports Exerc* 46:
1280 246, 2014.
- 1281 180. **Rock C, Newman V, Neuhouser M, Major J, Barnett M.** Antioxidant supplement use
1282 in cancer survivors and the general population. *J Nutr* 134: 3194S–3195S, 2004.
- 1283 181. **Rubin BB, Liauw S, Tittley J, Romaschin AD, Walker PM.** Prolonged adenine
1284 nucleotide resynthesis and reperfusion injury in postischemic skeletal muscle. *Am J Physiol*
1285 262: H1538–H1547, 1992.

- 1286 182. **Sachdev S, Davies KJ.** Production, detection, and adaptive responses to free radicals
1287 in exercise. *Free Radic Biol Med* 44: 215–223, 2008.
- 1288 183. **Saga N, Katamoto S, Naito H.** Effect of heat preconditioning by microwave
1289 hyperthermia on human skeletal muscle after eccentric exercise *J Sports Sci Med* 7: 176–
1290 183, 2008.
- 1291 184. **Samson L, Cairns J.** A new pathway for DNA repair in *Escherichia coli*. *Nature* 267:
1292 281–283, 1977.
- 1293 185. **Sandstrom ME, Zhang SJ, Bruton J, Silva JP, Reid MB, Westerblad H, Katz A.** Role of
1294 reactive oxygen species in contraction-mediated glucose transport in mouse skeletal
1295 muscle. *J Physiol* 575: 251–262, 2006.
- 1296 186. **Sapega AA, Heppenstall RB, Sokolow DP, Graham TJ, Maris JM, Ghosh AK, Chance
1297 B, Osterman AL.** The bioenergetics of preservation of limbs before replantation. The
1298 rationale for intermediate hypothermia. *J Bone Joint Surg Am* 70: 1500–1513, 1988.
- 1299 187. **Sato Y.** The history and future of Kaatsu training. *Int J Kaatsu Training Res* 1: 1–5,
1300 2005.
- 1301 188. **Schaser KD, Disch AC, Stover JF, Lauffer A, Bail HJ, Mittlmeier T.** Prolonged
1302 superficial local cryotherapy attenuates microcirculatory impairment, regional
1303 inflammation, and muscle necrosis after closed soft tissue injury in rats. *Am J Sports Med* 35:
1304 93–102, 2007.
- 1305 189. **Schaser KD, Stover JF, Melcher I, Lauffer A, Haas NP, Bail HJ, Stockle U, Puhl G,
1306 Mittlmeier TW.** Local cooling restores microcirculatory hemodynamics after closed soft-
1307 tissue trauma in rats. *J Trauma* 61: 642–649, 2006.
- 1308 190. **Schoenfeld BJ.** The use of nonsteroidal anti-inflammatory drugs for exercise-induced
1309 muscle damage: implications for skeletal muscle development. *Sports Med* 42: 1017–1028,
1310 2012.
- 1311 191. **Sekins KM, Lehmann JF, Esselman P, Dundore D, Emery AF, deLateur BJ, Nelp WB.**
1312 Local muscle blood flow and temperature responses to 915MHz diathermy as
1313 simultaneously measured and numerically predicted. *Arch Phys Med Rehabil* 65: 1–7, 1984.
- 1314 192. **Selsby JT, Dodd SL.** Heat treatment reduces oxidative stress and protects muscle
1315 mass during immobilization. *Am J Physiol Regul Integr Comp Physiol* 289: R134–R139, 2005.
- 1316 193. **Selsby JT, Rother S, Tsuda S, Prakash O, Quindry J, Dodd SL.** Intermittent
1317 hyperthermia enhances skeletal muscle regrowth and attenuates oxidative damage
1318 following reloading. *J Appl Physiol* 102: 1702–1707, 2007.
- 1319 194. **Selye H, Fortier C.** Adaptive reactions to stress. *Res Publ Assoc Res Nerv Ment Dis* 29:
1320 3–18, 1949.
- 1321 195. **Senf SM, Dodd SL, McClung JM, Judge AR.** Hsp70 overexpression inhibits NF-κB and
1322 Foxo3a transcriptional activities and prevents skeletal muscle atrophy. *FASEB J* 22: 3836–
1323 3845, 2008.
- 1324 196. **Senf SM, Howard TM, Ahn B, Ferreira LF, Judge AR.** Loss of the inducible Hsp70
1325 delays the inflammatory response to skeletal muscle injury and severely impairs muscle
1326 regeneration. *PLoS One* 8: e62687, 2013.
- 1327 197. **Serrano AL, Baeza-Raja B, Perdiguero E, Jardi M, Munoz-Canoves P.** Interleukin-6 is
1328 an essential regulator of satellite cell-mediated skeletal muscle hypertrophy. *Cell Metab* 7:
1329 33–44, 2008.
- 1330 198. **Shen W, Prisk V, Li Y, Foster W, Huard J.** Inhibited skeletal muscle healing in
1331 cyclooxygenase-2 gene-deficient mice: the role of PGE₂ and PGF_{2α}. *J Appl Physiol* 101: 1215–
1332 1221, 2006.

- 1333 199. **Shima Y, Kitaoka K, Yoshiki Y, Maruhashi Y, Tsuyama T, Tomita K.** Effect of heat
1334 shock preconditioning on ROS scavenging activity in rat skeletal muscle after downhill
1335 running. *J Physiol Sci* 58: 341–348, 2008.
- 1336 200. **Shinohara M, Kouzaki M, Yoshihisa T, Fukunaga T.** Efficacy of tourniquet ischemia
1337 for strength training with low resistance. *Eur J Appl Physiol Occup Physiol* 77: 189–191, 1998.
- 1338 201. **Silveira LR, Pilegaard H, Kusuhashi K, Curi R, Hellsten Y.** The contraction induced
1339 increase in gene expression of peroxisome proliferator-activated receptor (PPAR)- γ
1340 coactivator 1 α (PGC-1 α), mitochondrial uncoupling protein 3 (UCP3) and hexokinase II (HKII)
1341 in primary rat skeletal muscle cells is dependent on reactive oxygen species. *Biochim*
1342 *Biophys Acta* 1763: 969–976, 2006.
- 1343 202. **Smith WL, Meade EA, DeWitt DL.** Interactions of PGH synthase isozymes-1 and -2
1344 with NSAIDs. *Ann N Y Acad Sci* 744: 50–57, 1994.
- 1345 203. **Smuder AJ, Kavazis AN, Hudson MB, Nelson WB, Powers SK.** Oxidation enhances
1346 myofibrillar protein degradation via calpain and caspase-3. *Free Radic Biol Med* 49: 1152–
1347 1160, 2010.
- 1348 204. **Southam CM, Ehrlich J.** Effects of extracts of western red cedar heartwood on certain
1349 wood-decaying fungi in culture. *Phytopathology* 33: 517–524, 1943.
- 1350 205. **Spriet L.** New insights into the interaction of carbohydrate and fat metabolism during
1351 exercise. *Sports Med* 44: 87–96, 2014.
- 1352 206. **Steffen BT, Lees SJ, Booth FW.** Anti-TNF treatment reduces rat skeletal muscle
1353 wasting in monocrotaline-induced cardiac cachexia. *J Appl Physiol* 105: 1950–1958, 2008.
- 1354 207. **Strelkov AB, Fields AL, Baracos VE.** Effects of systemic inhibition of prostaglandin
1355 production on protein metabolism in tumor-bearing rats. *Am J Physiol* 257: C261–C269,
1356 1989.
- 1357 208. **Strobel NA, Peake JM, Matsumoto A, Marsh SA, Coombes JS, Wadley GD.**
1358 Antioxidant supplementation reduces skeletal muscle mitochondrial biogenesis. *Med Sci*
1359 *Sports Exerc* 43: 1017–1024, 2011.
- 1360 209. **Suga T, Okita K, Morita N, Yokota T, Hirabayashi K, Horiuchi M, Takada S, Omokawa**
1361 **M, Kinugawa S, Tsutsui H.** Dose effect on intramuscular metabolic stress during low-
1362 intensity resistance exercise with blood flow restriction. *J Appl Physiol* 108: 1563–1567,
1363 2010.
- 1364 210. **Suga T, Okita K, Morita N, Yokota T, Hirabayashi K, Horiuchi M, Takada S, Takahashi**
1365 **T, Omokawa M, Kinugawa S, Tsutsui H.** Intramuscular metabolism during low-intensity
1366 resistance exercise with blood flow restriction. *J Appl Physiol* 106: 1119–1124, 2009.
- 1367 211. **Suga T, Okita K, Takada S, Omokawa M, Kadoguchi T, Yokota T, Hirabayashi K,**
1368 **Takahashi M, Morita N, Horiuchi M, Kinugawa S, Tsutsui H.** Effect of multiple set on
1369 intramuscular metabolic stress during low-intensity resistance exercise with blood flow
1370 restriction. *Eur J Appl Physiol* 112: 3915–3920, 2012.
- 1371 212. **Summan M, Warren GL, Mercer RR, Chapman R, Hulderman T, Van Rooijen N,**
1372 **Simeonova PP.** Macrophages and skeletal muscle regeneration: a clodronate-containing
1373 liposome depletion study. *Am J Physiol Regul Integr Comp Physiol* 290: R1488–R1495, 2006.
- 1374 213. **Swenson C, Sward L, Karlsson J.** Cryotherapy in sports medicine. *Scand J Med Sci*
1375 *Sports* 6: 193–200, 1996.
- 1376 214. **Takagi R, Fujita N, Arakawa T, Kawada S, Ishii N, Miki A.** Influence of icing on muscle
1377 regeneration after crush injury to skeletal muscles in rats. *J Appl Physiol* 110: 382–388, 2011.

- 1378 215. **Takarada Y, Nakamura Y, Aruga S, Onda T, Miyazaki S, Ishii N.** Rapid increase in
1379 plasma growth hormone after low-intensity resistance exercise with vascular occlusion. *J*
1380 *Appl Physiol* 88: 61–65, 2000.
- 1381 216. **Takeuchi K, Hatade T, Wakamiya S, Fujita N, Arakawa T, Miki A.** Heat stress
1382 promotes skeletal muscle regeneration after crush injury in rats. *Acta Histochem* 116: 327–
1383 334, 2014.
- 1384 217. **Tamura Y, Matsunaga Y, Masuda H, Takahashi Y, Takahashi Y, Terada S, Hoshino D,**
1385 **Hatta H.** Post-exercise whole-body heat stress additively enhances endurance training-
1386 induced mitochondrial adaptations in mouse skeletal muscle. *Am J Physiol Regul Integr*
1387 *Comp Physiol* 307: R931–R943, 2014.
- 1388 218. **Tartibian B, Maleki B, Abbasi A.** Omega-3 fatty acids supplementation attenuates
1389 inflammatory markers after eccentric exercise in untrained men. *Clin J Sports Med* 21: 131–
1390 137, 2011.
- 1391 219. **Teixeira CF, Zamuner SR, Zuliani JP, Fernandes CM, Cruz-Hofling MA, Fernandes I,**
1392 **Chaves F, Gutierrez JM.** Neutrophils do not contribute to local tissue damage, but play a key
1393 role in skeletal muscle regeneration, in mice injected with Bothrops asper snake venom.
1394 *Muscle Nerve* 28: 449–459, 2003.
- 1395 220. **Theodorou AA, Nikolaidis MG, Paschalis V, Koutsias S, Panayiotou G, Fatouros IG,**
1396 **Koutedakis Y, Jamurtas AZ.** No effect of antioxidant supplementation on muscle
1397 performance and blood redox status adaptations to eccentric training. *Am J Clin Nutr* 93:
1398 1373–1383, 2011.
- 1399 221. **Tidball JG, Wehling-Henricks M.** Macrophages promote muscle membrane repair
1400 and muscle fibre growth and regeneration during modified muscle loading in mice in vivo. *J*
1401 *Physiol* 578: 327–336, 2007.
- 1402 222. **Tokmakidis SP, Kokkinidis EA, Smilios I, Douda H.** The effects of ibuprofen on
1403 delayed muscle soreness and muscular performance after eccentric exercise. *J Strength*
1404 *Cond Res* 17: 53–59, 2003.
- 1405 223. **Touchberry CD, Gupte AA, Bomhoff GL, Graham ZA, Geiger PC, Gallagher PM.** Acute
1406 heat stress prior to downhill running may enhance skeletal muscle remodeling. *Cell Stress*
1407 *Chaperones* 17: 693–705, 2012.
- 1408 224. **Trappe T, Carroll C, Dickinson J, LeMoine J, Haus J, Sullivan B, Lee J, Jemiolo B,**
1409 **Weinheimer E, Hollon C.** Influence of acetaminophen and ibuprofen on skeletal muscle
1410 adaptations to resistance exercise in older adults. *Am J Physiol Regul Integr Comp Physiol*
1411 300: R655–R662, 2011.
- 1412 225. **Trappe T, Raue U, Williams R, Carrithers J, Hickner R.** Effects of age and resistance
1413 exercise on skeletal muscle interstitial prostaglandin $F_{2\alpha}$. *Prostaglandins Leukot Essent Fatty*
1414 *Acid* 74: 175–181, 2006.
- 1415 226. **Trappe T, Standley R, Jemiolo B, Carroll C, Trappe S.** Prostaglandin and myokine
1416 involvement in the cyclooxygenase-inhibiting drug enhancement of skeletal muscle
1417 adaptations to resistance exercise in older adults. *Am J Physiol Regul Integr Comp Physiol*
1418 304: R198–R205, 2013.
- 1419 227. **Trappe TA, Fluckey JD, White F, Lambert CP, Evans WJ.** Skeletal muscle $PGF_{2\alpha}$ and
1420 PGE_2 in response to eccentric resistance exercise: influence of ibuprofen acetaminophen. *J*
1421 *Clin Endocrinol Metab* 86: 5067–5070, 2001.
- 1422 228. **Trappe TA, Liu SZ.** Effects of prostaglandins and COX-inhibiting drugs on skeletal
1423 muscle adaptations to exercise. *J Appl Physiol* 115: 909–919, 2013.

- 1424 229. **Trappe TA, White F, Lambert CP, Cesar D, Hellerstein M, Evans WJ.** Effect of
 1425 ibuprofen and acetaminophen on postexercise muscle protein synthesis. *Am J Physiol*
 1426 *Endocrinol Metab* 282: E551–E556, 2002.
- 1427 230. **Vaile J, Halson S, Gill N, Dawson B.** Effect of hydrotherapy on recovery from fatigue.
 1428 *Int J Sports Med* 29: 539–544, 2008.
- 1429 231. **Vaile J, Halson S, Gill N, Dawson B.** Effect of hydrotherapy on the signs and
 1430 symptoms of delayed onset muscle soreness. *Eur J Appl Physiol* 102: 447–455, 2008.
- 1431 232. **Vane JR.** Inhibition of prostaglandin synthesis as a mechanism of action for aspirin-
 1432 like drugs. *Nat New Biol* 231: 232–235, 1971.
- 1433 233. **Vella L, Markworth JF, Peake JM, Snow RJ, Cameron-Smith D, Russell AP.** Ibuprofen
 1434 supplementation and its effects on NF- κ B activation in skeletal muscle following resistance
 1435 exercise. *Physiol Rep* 2: 2014.
- 1436 234. **Venditti P, Napolitano G, Barone D, Di Meo S.** Vitamin E supplementation modifies
 1437 adaptive responses to training in rat skeletal muscle. *Free Radic Res* 48: 1179–1189, 2014.
- 1438 235. **Veskoukis AS, Nikolaidis MG, Kyparos A, Kokkinos D, Nepka C, Barbanis S, Kouretas**
 1439 **D.** Effects of xanthine oxidase inhibition on oxidative stress and swimming performance in
 1440 rats. *Appl Physiol Nutr Metab* 33: 1140–1154, 2008.
- 1441 236. **Vollaard NB, Shearman JP, Cooper CE.** Exercise-induced oxidative stress: myths,
 1442 realities and physiological relevance. *Sports Med* 35: 1045–1062, 2005.
- 1443 237. **Wadley GD, McConell GK.** High-dose antioxidant vitamin C supplementation does
 1444 not prevent acute exercise-induced increases in markers of skeletal muscle mitochondrial
 1445 biogenesis in rats. *J Appl Physiol* 108: 1719–1726, 2010.
- 1446 238. **Wadley GD, Nicolas MA, Hiam D, McConell GK.** Xanthine oxidase inhibition
 1447 attenuates skeletal muscle signaling following acute exercise but does not impair
 1448 mitochondrial adaptations to endurance training. *Am J Physiol Endocrinol Metab* 304: E853–
 1449 E862, 2013.
- 1450 239. **Wernbom M, Apro W, Paulsen G, Nilsen TS, Blomstrand E, Raastad T.** Acute low-
 1451 load resistance exercise with and without blood flow restriction increased protein signalling
 1452 and number of satellite cells in human skeletal muscle. *Eur J Appl Physiol* 113: 2953–2965,
 1453 2013.
- 1454 240. **Wernbom M, Augustsson J, Raastad T.** Ischemic strength training: a low-load
 1455 alternative to heavy resistance exercise? *Scand J Med Sci Sports* 18: 401–416, 2008.
- 1456 241. **Wernbom M, Paulsen G, Nilsen TS, Hisdal J, Raastad T.** Contractile function and
 1457 sarcolemmal permeability after acute low-load resistance exercise with blood flow
 1458 restriction. *Eur J Appl Physiol* 112: 2051–2063, 2012.
- 1459 242. **Wojtaszewski J, MacDonald C, Nielsen J, Hellsten Y, Hardie D, Kemp B, Kiens B,**
 1460 **Richter E.** Regulation of 5'AMP-activated protein kinase activity and substrate utilization in
 1461 exercising human skeletal muscle. *Am J Physiol Endocrinol Metab* 284: E813–E822, 2003.
- 1462 243. **Wright DC, Han DH, Garcia-Roves PM, Geiger PC, Jones TE, Holloszy JO.** Exercise-
 1463 induced mitochondrial biogenesis begins before the increase in muscle PGC-1 α expression. *J*
 1464 *Biol Chem* 282: 194–199, 2007.
- 1465 244. **Wu Z, Puigserver P, Andersson U, Zhang C, Adelmant G, Mootha V, Troy A, Cinti S,**
 1466 **Lowell B, Scarpulla RC, Spiegelman BM.** Mechanisms controlling mitochondrial biogenesis
 1467 and respiration through the thermogenic coactivator PGC-1. *Cell* 98: 115–124, 1999.
- 1468 245. **Wyper DJ, McNiven DR.** The effect of microwave therapy upon muscle blood flow in
 1469 man. *Br J Sports Med* 10: 19–21, 1976.

- 1470 246. **Yamane M, Teruya H, Nakano M, Ogai R, Ohnishi N, Kosaka M.** Post-exercise leg and
 1471 forearm flexor muscle cooling in humans attenuates endurance and resistance training
 1472 effects on muscle performance and on circulatory adaptation. *Eur J Appl Physiol* 96: 572–
 1473 580, 2006.
- 1474 247. **Yamashita-Goto K, Ohira Y, Okuyama R, Sugiyama H, Honda M, Sugiura T, Yamada**
 1475 **S, Akema T, Yoshioka T.** Heat stress facilitates stretch-induced hypertrophy of cultured
 1476 muscle cells. *J Gravit Physiol* 9: P145–146, 2002.
- 1477 248. **Yellon DM, Latchman DS, Marber MS.** Stress proteins—an endogenous route to
 1478 myocardial protection: fact or fiction? *Cardiovasc Res* 27: 158–161, 1993.
- 1479 249. **Yeo W, McGee S, Carey A, Paton C, Garnham A, Hargreaves M, Hawley J.** Acute
 1480 signalling responses to intense endurance training commenced with low or normal muscle
 1481 glycogen. *Exp Physiol* 95: 351–358, 2010.
- 1482 250. **Yeo W, Paton C, Garnham A, Burke L, Carey A, Hawley J.** Skeletal muscle adaptation
 1483 and performance responses to once a day versus twice every second day endurance training
 1484 regimens. *J Appl Physiol* 105: 1462–1470, 2008.
- 1485 251. **Yfanti C, Akerstrom T, Nielsen S, Nielsen AR, Mounier R, Mortensen OH, Lykkesfeldt**
 1486 **J, Rose AJ, Fischer CP, Pedersen BK.** Antioxidant supplementation does not alter endurance
 1487 training adaptation. *Med Sci Sports Exerc* 42: 1388–1395, 2010.
- 1488 252. **Yfanti C, Fischer CP, Nielsen S, Akerstrom T, Nielsen AR, Veskoukis AS, Kouretas D,**
 1489 **Lykkesfeldt J, Pilegaard H, Pedersen BK.** Role of vitamin C and E supplementation on IL-6 in
 1490 response to training. *J Appl Physiol* 112: 990–1000, 2012.
- 1491 253. **Yfanti C, Nielsen AR, Akerstrom T, Nielsen S, Rose AJ, Richter EA, Lykkesfeldt J,**
 1492 **Fischer CP, Pedersen BK.** Effect of antioxidant supplementation on insulin sensitivity in
 1493 response to endurance exercise training. *Am J Physiol Endocrinol Metab* 300: E761–E770,
 1494 2011.
- 1495 254. **Zhang SJ, Sandstrom ME, Lanner JT, Thorell A, Westerblad H, Katz A.** Activation of
 1496 aconitase in mouse fast-twitch skeletal muscle during contraction-mediated oxidative stress.
 1497 *Am J Physiol Cell Physiol* 293: C1154–C1159, 2007.
- 1498 255. **Ørtenblad N, Nielsen J, Saltin B, Holmberg H-C.** Role of glycogen availability in
 1499 sarcoplasmic reticulum Ca^{2+} kinetics in human skeletal muscle. *J Physiol* 589: 711–725, 2011.
- 1500 256. **Ørtenblad N, Westerblad H, Nielsen J.** Muscle glycogen stores and fatigue. *J Physiol*
 1501 591: 4405–4413, 2013.

1502 **Tables****Table 1.** Effects of glycogen concentration on physiological responses to exercise in human skeletal muscle.

Reference	Design	$\Delta\%$	Low glycogen	Findings
(242)	Acute	-82%	$163 \text{ mmol} \cdot \text{kg}^{-1} \cdot \text{dw}$	\uparrow AMPK activity
(9)	Acute	-75%	$103 \text{ mmol} \cdot \text{kg}^{-1} \cdot \text{dw}$	\uparrow p53 phosphorylation \uparrow Mitochondrial mRNA
(172)	Acute	-65%	$166 \text{ mmol} \cdot \text{kg}^{-1} \cdot \text{dw}$	\uparrow Mitochondrial mRNA
(13)	Acute	-47%	$167 \text{ mmol} \cdot \text{kg}^{-1} \cdot \text{dw}$	\uparrow Protein degradation
(72)	Acute	-30%	$290 \text{ mmol} \cdot \text{kg}^{-1} \cdot \text{dw}$	\uparrow Leucine oxidation \downarrow Net protein balance
(23)	Acute	-52%	$180 \text{ mmol} \cdot \text{kg}^{-1} \cdot \text{dw}$	\leftrightarrow Muscle protein synthesis
(255)	Acute	-69%	$167 \text{ mmol} \cdot \text{kg}^{-1} \cdot \text{dw}$	\downarrow SR Ca^{2+} release rate
(50)	Acute	-68%	$245 \text{ mmol} \cdot \text{kg}^{-1} \cdot \text{dw}$	\downarrow SR Ca^{2+} release rate
(65)	Chronic	-68%	$210 \text{ mmol} \cdot \text{kg}^{-1}$	\uparrow Mitochondrial enzyme activity
(250)	Chronic	-50%	$250 \mu\text{mol} \cdot \text{g}^{-1} \cdot \text{dw}$	\uparrow Mitochondrial enzymes \uparrow Fat oxidation

dw, dry weight; SR, sarcoplasmic reticulum.

1503

1504

Table 2. Summary of studies investigating the effects of heat stress on muscle regeneration.

Reference	Study type	Treatment	Assessment period	Outcome variables
(48)	Rats; ischemia	Hot water @ 42.5°C Duration: 20 min Timing: 12 h preinjury	1.5 h postinjury	Electron microscopy, PCr, ATP, HSP72
(199)	Rats; downhill running	Heat chamber at 42°C Duration: 60 min Timing: 48 h preinjury	1, 2, 3, and 7 d postinjury	ROS production and scavenging, HSP72, histology
(223)	Rats; downhill running	Hot water @ 43°C; Duration: 20 min Timing: 48 h preinjury	2 h and 2 d postinjury	Histology, Akt, p70S6K, ERK1/2, JNK, HSP72, HSP25, MHC
(96)	Rats; cardiotoxin injury	Heat chamber at 41°C Duration: 60 min Timing: 24 h preinjury or 0 h postinjury	1, 3, 7, 14, and 28 d postinjury	Muscle mass, central nucleated fibers, fiber CSA, HSP72, Pax7
(25)	Rats; acute strain injury	Infrared lamp Duration: 5 min Timing: 30 min and 2×/day postinjury	1, 5, 10, and 15 d postinjury	Lipid peroxidation, antioxidant enzymes myeloperoxidase
(154)	Rats; cardiotoxin injury	Hot water @ 42°C Duration: 30 min Timing: 48 h postinjury and then every second day	7 and 15 d postinjury	Fiber CSA, myonuclei, Pax7, M-Cadherin, MyoD, HSP72, calcineurin
(71)	Rats; tenotomy	Heat chamber @ 40.5–41°C Duration: 30 min Timing: 24 h preinjury; 1–6 d postinjury	7 d postinjury	Muscle mass, histology, fiber CSA, HSP72, collagen, TGF-β1, MMP-2, MMP-9, TIMP
(216)	Rats; acute crush injury	Hot pack @ 42°C Duration: 20 min Timing: 5 min postinjury	6 and 12 h; 1–7, 14, and 28 d postinjury	Central nucleated fibers Fiber CSA, macrophages TGF-β1, IGF-1, Pax7, collagen
(68)	Rats; acute crush injury	Hot pack @ 42°C; Duration: 20 min Timing: 5 min postinjury	12 h; 1–5, 7, 14, and 28 d postinjury	MyoD, myogenin, PCNA Pax7

(217)	Mice; acute treadmill running	Heat chamber @ 41°C Duration: 30 min Timing: Immediately postexercise	30 min postexercise	AMPK, ACC, p38 MAPK, CaMKII, Akt, mTOR p70S6K
-------	--	---	------------------------	--

Abbreviations: PCr, phosphocreatine; MHC, myosin heavy chain; PCNA, proliferating cells nuclear antigen; ROS, reactive oxygen species; CaMK, calmodulin-dependent protein kinase; CSA, cross-sectional area; Akt, protein kinase B; MMP, matrix metalloproteinase; TIMP, tissue inhibitor of matrix metalloproteinase. See Figure 1 for details of other abbreviations.

1506

1507

Table 3. Studies investigating the effects of cryotherapy on muscle regeneration.				
Reference	Study type	Treatment	Assessment period	Outcome variables
(214)	Rat; acute crush injury	Topical icing Duration: 20 min Timing: 5 min postinjury	6 and 12 h; 1–7, 14, and 28 d postinjury	Central nucleated fibers Fiber CSA, macrophage , TGF- β 1, IGF-1, Pax7, collagen
(25)	Rat; acute crush injury	Topical icing Duration: 5 min Timing: 30 min and 2 \times /d postinjury	1, 5, 10, and 15 d postinjury	Lipid peroxidation Antioxidant enzymes Myeloperoxidase
(174)	Rat; acute contusion injury	Topical icing Duration: 5 min Timing: Immediately and 6 h postinjury	1 d postinjury	Lipid peroxidation Antioxidant enzymes Myeloperoxidase Na ⁺ -K ⁺ ATPase, Ca ²⁺ ATPase Lactate dehydrogenase
(76)	Rat; acute contusion injury	Topical icing Duration: 5 min; intermittently for 1 h Timing: Immediately postinjury or 24 h postinjury	1, 2, and 6 h; 1, 2, 5, and 7 d postinjury	Neutrophil infiltration Macrophage infiltration Desmin ⁺ myoblasts
(102)	Rat; acute contusion injury	Cold saline (3°C) infusion Duration: 10 min Timing: 5 min postinjury	15 min postinjury	Leukocyte rolling and adhesion
(189)	Rat; acute contusion injury	Cold saline (8°C) infusion Duration: 20 min Timing: ~20 min postinjury	1 h postinjury	Edema, microvascular perfusion, leukocyte rolling/adhesion Neutrophils and macrophages
(188)	Rat; acute contusion injury	Cold saline (8°C) infusion Duration: 6 h Timing: ~20 min postinjury	1 d postinjury	Edema, microvascular perfusion, leukocyte rolling/adhesion Neutrophils and macrophages Desmin expression

CSA, cross-sectional area; TGF, transforming growth factor

1508

Table 4. Summary of physiological and molecular responses, acute and chronic adaptations to treatments that enhance or dampen exercise-induced hormesis in skeletal muscle.

	Treatments that dampen hormesis				Treatments that enhance hormesis			
	Cryotherapy	NSAIDs	Antioxidant supplementation		Carbohydrate restriction		Heat stress	Blood flow restriction
Physiological rationale	Analgesia ↓ Muscle blood flow ↓ Inflammation ↑ Hydrostatic pressure	Analgesia ↓ Inflammation	↓ Oxidative stress		↑ Metabolic stress		↓ Muscle breakdown	↑ Metabolic stress ↑ Oxidative stress ↑ Blood pooling
Cells and signalling molecules upregulated	TGF-β	IL-6 MCP-1 Cyclooxygenase 2			AMPK ACC p53 PGC-1α CS	SDH HAD COXIV PDK4	Macrophages CS HSPs MyoD p38 MAPK p70S6K Myogenin mTOR	Pax7 AMPK MAPK HSPs
Cells and signalling molecules downregulated	Neutrophils Macrophages IGF-1 Pax7	Prostaglandins ERK/RSK/MNK p70S6K/rpS6 Leukotrienes Resolving mediators	p38 MAPK ERK AMPK IL-6 NFκB	PGC-1α Tfam COX SOD			Macrophages NFκB AMPK ACC	
Acute effects	↓ Soreness	Soreness? ↔ Inflammation ↓ Protein synthesis ↓ Satellite cells			↓ SR Ca ²⁺ release rate ↑ Protein breakdown	↓ Loss of strength* ↓ Soreness* ↓ Swelling ↑ Range of motion*		↑ Loss of strength ↑ Soreness ↑ Swelling
Chronic effects	↓ Fibre CSA ↑ Fibrosis ↓ Strength	Young healthy ↔ muscle mass?	↓ Antioxidant enzymes		↑ Mitochondrial enzymes ↑ Fat oxidation	↑ Mitochondrial enzymes ↑ Respiratory chain protein content		↑ Hypertrophy

↔ strength?

↔ Performance

Elderly

↑ muscle mass

↑ strength

Abbreviations: TGF, transforming growth factor; MCP, monocyte chemotactic protein; AMPK, adenosine monophosphate activated protein kinase; ACC, acetyl-CoA-carboxylase; PGC, peroxisome proliferator-activated receptor coactivator; CS, citrate synthase; SDH, succinate dehydrogenase; HAD, hydroxyacyl-CoA-dehydrogenase; COX, cytochrome oxidase; PDK, pyruvate dehydrogenase kinase; HSP, heat shock protein; Pax, paired box protein; mTOR; mammalian target of rapamycin; Mnk, MAPK-interacting kinase; RSK, p90 ribosomal S6 kinase; rpS6, ribosomal S6 kinase; Tfam, mitochondrial transcription factor A; SOD, superoxide dismutase; CSA, cross-sectional area. ↔ no change. * conflicting evidence for an increase/decrease or no change.